



**THE MARINE HABITAT CLASSIFICATION
FOR BRITAIN AND IRELAND
VERSION 04.05**

INTRODUCTION

**DAVID W. CONNOR, JAMES H. ALLEN*, NEIL GOLDING, KERRY L.
HOWELL, LOUISE M. LIEBERKNECHT, KATE O. NORTHEN[†] AND
JOHNNY B. REKER**

May 2004



JOINT NATURE CONSERVATION COMMITTEE

*** INSTITUTE OF COASTAL AND ESTUARINE STUDIES, UNIVERSITY OF HULL**

† SEA-SCOPE, DEVON

Joint Nature Conservation Committee
Monkstone House
City Road
Peterborough PE1 1JY
UK

Recommended citation:

DAVID W. CONNOR, JAMES H. ALLEN, NEIL GOLDING, KERRY L. HOWELL, LOUISE M. LIEBERKNECHT, KATE O. NORTHEN AND JOHNNY B. REKER (2004)

The Marine Habitat Classification for Britain and Ireland Version 04.05.

JNCC, Peterborough ISBN 1 861 07561 8 (internet version)
www.jncc.gov.uk/MarineHabitatClassification

ISBN 1 861 07561 8

Text © 2004 Copyright JNCC All Rights Reserved

Contents

Summary	5
Acknowledgements	6
Rationale and need for a habitat classification	8
Purpose and scope	8
Requirements of a habitat classification system.....	8
Applications	9
Nature of the marine environment	11
The habitat scale in characterising the marine environment	11
Environmental influences at the habitat scale	11
Terminology: the terms biotope, habitat and community.....	12
Classification development - approach and methods used	16
Review of classification systems and literature.....	16
Consultation and testing	16
Field surveys and other data acquisition	16
Data analysis	17
Littoral rock.....	17
Littoral sediment	17
Infralittoral rock	19
Circalittoral rock	19
Sublittoral sediment.....	19
Structure of the classification	21
A framework for the classification (EUNIS levels 2 and 3).....	21
Development of a hierarchical classification	25
Top-down classification	25
Bottom-up classification.....	25
Distinguishing and defining types	27
How to use the classification	29
Habitat matrices.....	29
Hierarchy structure diagrams	29
Hierarchical list of types.....	29
Layout of descriptions for each type	29
Comparative tables	31
Species nomenclature	32
Understanding the codes	32

Field recording and data management	43
Terms used for field recording and habitat definition	43
MNCR SACFOR abundance scales	46
References	48

Summary

This publication presents a fully revised version of the national classification of benthic marine habitats (seashore and seabed habitats and their associated communities of species) for Britain and Ireland. It was originally developed by JNCC's Marine Nature Conservation Review (MNCR) as part of the EC Life Nature-funded BioMar project (Connor *et al.* 1997 a, b).

The classification provides a tool to aid the management and conservation of marine habitats. It has been developed through the analysis of empirical data sets, the review of other classifications and scientific literature, and in collaboration with a wide range of marine scientists and conservation managers. It is fully compatible with and contributes to the European EUNIS habitat classification system (<http://mrw.wallonie.be/dgrne/sibw/EUNIS/home.html>).

An outline of the rationale, uses, overall structure and development methods is given in this introductory document. The classification website (www.jncc.gov.uk/MarineHabitatClassification) contains a full listing of newly defined and revised classification types, a detailed description of each type, distribution maps and, where possible, colour photographs. Separate documents are available to download on the website, containing biotope descriptions for each of the sections of the classification.

The classification is presented in hierarchical format, and through a series of habitat matrices. It comprises:

		Number of types defined
Level 1	Environment (marine)	1
Level 2	Broad habitat types	5
Level 3	Habitat complexes	24
Level 4	Biotope complexes	75
Levels 5 & 6	Biotoxes and sub-biotoxes	370

IMPORTANT

This classification supersedes versions 97.06 and 03.02.

Users of the classification must ensure they state which version has been used in any reports, data interpretation or field survey.

Acknowledgements

The development and success of the classification has only been possible through the considerable input and tremendous enthusiasm of a wide variety of people. Expertise from scientific and conservation management perspectives, with international through to local standpoints, and with views on both general philosophies and practical considerations have been essential to ensure the classification has developed as a robust scientific but practical tool for marine nature conservation and environmental management. We are very grateful to all those involved, for both the many positive comments which have encouraged us and the criticisms which have helped to sharpen the end product.

The following have particularly contributed to the present version:

Analysis of data:

Kate Bull (JNCC, now EN)
Rohan Holt (CCW)
Christine Howson (Consultant)
Eammon Kelly (JNCC)
Kate Smith (JNCC, now CCW)

Comments on the revised classification:

Kathryn Baukham (CCW)
John Baxter (SNH)
Daryl Birkett (Queen's University, Belfast)
Martin Bradley (EHS)
Paul Brazier (CCW)
Alison Brunstrom (CCW)
Francis Bunker (Marine Seen)
Ann Bunker (CCW)
Paul Gilliland (EN)
Mike Guiry (National University of Ireland, Galway)
Eleanor Hill (EN)
Tim Hill (EN)
Keith Hiscock (Marine Biological Association)
Monica Jones (CCW)
Charles Lindenbaum (CCW)
Natasha Lough (CCW)
Julia Nunn (Ulster Museum)
Brendan O'Connor (Aquafact)
Ivor Rees (University of Wales, Bangor)
Kate Smith (CCW)
Christina Vina-Herbon (North Atlantic Fisheries College)
Tim Worsfold (Unicomarine)
Nicola White (Posford Haskoning)

Data management and development of the electronic publication:

Dave Chambers
Mike Webster
Steve Wilkinson
Ulric Wilson
Hannah Betts
Cathy Gardner

Introduction

This publication presents a classification of marine habitats for the shores and seabed around Britain and Ireland. It has been substantially revised and updated from the previous version which was developed by JNCC's Marine Nature Conservation Review (MNCR) as part of the EC Life Nature-funded BioMar programme (Connor *et al.* 1997a, b). This new version reflects the continuing needs of end-users for improved information about each habitat, which is presented in a variety of interactive ways in this web-based version. As a result of re-analysis of field data, including new data from recent surveys in candidate Special Areas of Conservation (cSACs), a number of new habitat types have been described. The classification has been restructured, both to reflect improved understanding of the inter-relationship of habitat types and to align itself more closely with the European EUNIS habitat classification system (<http://mrw.wallonie.be/dgrne/sibw/EUNIS/home.html>).

Rationale and need for a habitat classification

Purpose and scope

Seabed habitats and the communities of species that occupy them are an essential component of the marine ecosystem and our overall understanding of ecosystem function must relate seabed habitats to hydrography, nutrient cycling, plankton changes and the distribution of wide-ranging species (i.e. fish stocks, marine mammals, birds). A greater understanding of the distribution, extent and status or quality of marine habitats is required to facilitate the protection of threatened and rare habitats and, more generally, the assessment of the state of the marine environment. Such information is also needed to improve spatial and strategic planning of human activities, in particular to promote the wiser use of habitats where there are competing demands (e.g. fishing, sand and gravel extraction, wind energy generation, nature conservation). As such, information on marine habitats needs to play a major role in the ecosystem-based approach to management of the marine environment that is now widely advocated at national and international levels (Defra 2002; North Sea Conference 2002).

This habitat classification has, consequently, been developed as a tool to aid the management and conservation of marine habitats. It provides an ecologically-based classification of seashore and seabed features, aimed primarily at classifying benthic communities of invertebrates and seaweeds in a way which is meaningful both to detailed scientific application and to the much broader requirements for management of the marine environment. The classification is relevant to the habitat requirements of more mobile species, such as fish and marine mammals, but these are not its primary focus. Whilst the corresponding European EUNIS classification also includes water column (plankton) habitats, this aspect has not yet been developed here.

The classification aims to provide comprehensive coverage, by including habitats for artificial, polluted or barren areas as well as more natural habitats, which encompass:

1. **Marine, estuarine and brackish-water (lagoon) habitats** - It also includes reference to saltmarsh habitats described in the National Vegetation Classification (NVC) (Rodwell 2000; Doody, Johnston & Smith 1993) as these are regularly covered by the sea, and NVC types which occur in brackish lagoons (Rodwell 1995).
2. **Rock and sediment habitats.**
3. **Upper shore to coastal waters** - From the supralittoral or splash zone and strand-line on the shore out to the 200 nm limit. The habitats beyond the near-shore subtidal zone (about the 3 mile/5 km limit) and below about 50 m depth are less well described here, due to more limited availability of data; more types will be defined as data become available.
4. **Plant and animal communities, including epibiota and infauna** - Types are defined using both their fauna and flora. Most benthic marine habitats include sedentary animals and small mobile animals which are an integral part of the community, whilst in many habitats, especially in deeper water, there are no plants (seaweeds or marine angiosperms) to characterise the habitats. Sediment types are defined both by their epibiota (surface-dwelling animals and plants) and their infauna (animals living in the sediment).
5. **Britain and Ireland** - It covers habitats throughout Britain and Ireland and, through a widely-accepted broad framework, is readily expandable to include offshore continental shelf habitats and other areas in the north-east Atlantic, Mediterranean and Baltic Seas. This is being achieved through the EUNIS classification.

Requirements of a habitat classification system

To underpin management and conservation of the marine environment, a habitat classification system should:

- be scientifically sound, adopting a logical structure in which the types are clearly defined on ecological grounds, avoiding overlap in their definition and duplication of types in different

parts of the system, and ensuring that ecologically-similar types are placed near to each other and at an appropriate level (within a hierarchical classification);

- provide a common and easily understood language for the description of marine habitats;
- be comprehensive, accounting for all the marine habitats within its geographic scope;
- be practical in format and clear in its presentation;
- include sufficient detail to be of practical use for conservation managers and field surveyors but be sufficiently broad (through hierarchical structuring) to enable summary habitat information to be presented at national and international levels or its use by non-specialists;
- be sufficiently flexible to enable modification resulting from the addition of new information, but stable enough to support ongoing uses. Changes should be clearly documented to enable reference back to previous versions (where possible, newly defined types need to be related back to types in earlier versions of the classification).

The following considerations were taken into account in establishing the classification:

- its intended application by a variety of users and at various scales (environmental managers, marine scientists and field surveyors working at local, national and international levels);
- the variety of intended applications;
- the variation in the scale of physical and biological features (recognising that marine ecosystems operate at a wide variety of scales, e.g. whole estuaries, individual mussel beds);
- the different levels of detail in available data;
- the different skill levels of future users and their different methods of survey.

Applications

A number of applications for the habitat classification system have been identified:

- to provide a practical system for the consistent description of habitat types;
- to map habitats to assess their geographical distribution;
- to map habitats to assess their extent;
- to provide categories for the assessment of the state of marine biological communities;
- to assess changes in habitat distribution and extent over time, to provide information on quality status, and rate of change in habitat distribution;
- to assess the relative importance of particular habitats (i.e. which habitats are rare or of national or regional importance) and the implications of this for prioritising management and conservation action. Such assessment can lead to the listing of habitats for conservation action (e.g. Red lists);
- to enable the nature conservation value of habitats at specific sites to be assessed, such as in the identification of marine protected areas (MPAs);
- to enable an assessment of the extent of protection afforded to habitats by existing or proposed MPAs and the degree to which this provides sufficient protection;
- to enable the range and intensity of human activities that occur in particular habitats, and the degree to which such habitats are affected by those activities, to be systematically assessed;
- to facilitate presentation of habitat information at a scale and level of detail that enables appropriate management action to be taken. Such presentation should be flexible to address a variety of biodiversity and management issues;

- habitat mapping information needs to be used in conjunction with other spatial information in Geographical Information Systems (GIS), particularly activities, management and conservation areas, and other environmental data sets.

Nature of the marine environment

The habitat scale in characterising the marine environment

The marine environment can be described or characterised at a number of different scales, ranging from ocean-level processes through to those that occur at species and genetic level (Connor *et al.* 2002). The scales of relevance here are marine landscapes, habitats and species; their inter-relationship can be expressed as follows:

- **Species** provide the globally accepted original classification of biological diversity, with well-established rules of taxonomy to distinguish between different types. Their classification is arranged in a hierarchy of genera, families, orders, classes and phyla.
- **Habitats** comprise suites of species (communities or assemblages) that consistently occur together, but which are derived from different parts of the taxonomic hierarchy (e.g. kelps, molluscs and fish in a kelp forest habitat). Their classification can also be structured in a hierarchy (biotopes, biotope complexes, broad habitats), reflecting degrees of similarity.
- **Marine Landscapes** comprise suites of habitats that consistently occur together, but which are often derived from different parts of the habitat classification hierarchy (e.g. saltmarsh, intertidal mudflats, rocky shores and subtidal mussel beds in an estuary).

The approach to classification or characterisation at each scale differs, each adopting differing factors to suit the requirements at that scale. Whilst the classification (taxonomy) of species, and to a lesser degree habitats, is now well established the seascape concept and their characterisation is a more recent approach to characterisation of the marine environment (Laffoley *et al.* 2000, Day & Roff 2000). The marine landscape concept was applied to the seabed and water column of the Irish Sea as part of the Irish Sea Pilot project (www.jncc.gov.uk/IrishSeaPilot).

Environmental influences at the habitat scale

Each species tends to live within a certain environment; that is, it has a preference for a combination of environmental factors (a niche), such as the substratum, temperature, salinity and hydrodynamic conditions that it is able to live within. The tolerance to different environmental conditions varies between species; it can be rather broad for some very common species but much more tightly defined for others. The niche occupied by a species may vary both temporally and spatially and is influenced not only by its physiological requirements and tolerance to change but also by the interactions between species, i.e. competition and predator-prey relationships.

In any particular place on the shore or seabed, a suite of species will occur, each adapted to the particular environmental conditions of that place, such as the conditions of an intertidal mudflat. Where such a suite of species occurs in other locations under similar environmental conditions, it can be defined as a **community** (or association or assemblage) of species which is occurring within a particular **habitat** type. The collective term **biotope** is now in common usage to encompass both of these biotic and abiotic elements.

Shore and seabed habitats are colonised primarily by seaweeds (on the shore and in shallow water) and by marine invertebrates from a wide range of phyla. Lichens (in the splash zone), higher plants (especially in saltmarshes) and fish contribute to a lesser degree. In contrast to terrestrial habitats, it is commonplace for marine habitats to be characterised, i.e. dominated, by animals rather than plants, and for the substratum to provide the main structure to the habitat (rather than plants such as in a forest).

Only a proportion of habitats have obvious dominant species (e.g. kelp forests, mussel beds, maerl beds). Many, particularly in deeper water, support a mosaic of species, none of which is particularly dominant, which may exhibit a degree of patchiness over the seashore or seabed and, in some cases, vary markedly with time. In these respects the species offer a much less robust mechanism for structuring a classification system than does the physical habitat in which they occur.

In the marine environment, there is a strong relationship between the abiotic nature of the habitat and the biological composition of the community it supports. Most communities appear to occur within a recognisable suite of environmental factors, although some occur within a more tightly-defined set of factors (habitat). One of the most important factors influencing species composition is the type of substratum present, which can be broadly divided into rock and sediment (the latter is closely linked to the hydrodynamic regime) whilst in estuaries salinity is an important factor. Community structure is additionally modified by biological factors such as recruitment, predation, grazing and inter-species competition. Species may modify habitats by their boring, accretion and bioturbation. The most important habitat attributes which appear to influence community composition are described in Table 1. In addition to habitat factors, biological and anthropogenic influences affect community composition. Some aspects of anthropogenic influence are outlined in Table 2.

Terminology: the terms biotope, habitat and community

A **biotope** is defined as the combination of an abiotic habitat and its associated community of species. It can be defined at a variety of scales (with related corresponding degrees of similarity) and should be a regularly occurring association to justify its inclusion within a classification system.

A **habitat** is taken to encompass the substratum (rock, sediment or biogenic reefs such as mussels), its topography and the particular conditions of wave exposure, salinity, tidal currents and other water quality characteristics (e.g. turbidity and oxygenation) which contribute to the overall nature of a place on the shore or seabed.

The term **community** is used here to mean an association of species which has particular species, at certain densities, in common.

Although communities are influenced by biological interactions (e.g. predation, recruitment processes) and by interference from certain human activities, their overall character is very strongly determined by the nature of the surrounding abiotic conditions. This consistent relationship between the biotic and abiotic elements is fundamental to the structure of the classification system. Types can be defined at a variety of scales, enabling the development of a hierarchical classification of types. The degree of similarity varies depending upon the scale considered, with more broadly defined types (e.g. sheltered rocky shores) having a lower level of similarity compared with more finely defined types (e.g. a lower shore sheltered rocky biotope).

Whilst the term habitat, as used here, is its more accepted scientific meaning, the term is more widely used, for instance in the EC Habitats Directive, to also include the community of species living in the habitat; the common use of the term is, therefore, synonymous with the term biotope.

Table 1 Environmental factors which influence community structure

<i>Factor</i>	<i>Rocky habitats</i>	<i>Sediment habitats</i>
<i>Substratum</i>	Varies from bedrock, through boulders to stony plains, often mixed with sediment. The degree of stability of the rock is important, with algae and animals increasingly able to colonise smaller stones in more sheltered stable conditions.	Ranges from shingle (mobile cobbles and pebbles), through gravel and sand to very soft mud and muddy gravels. The type of sediment, mainly determined by the dynamics of water movement at the site, is highly important in structuring community composition, although salinity may become more critical in upper estuarine conditions.
<i>Zonation: emersion / immersion on the shore (desiccation); depth in the subtidal (illumination) (see Figure 1 and Table 5 for further details)</i>	A major factor, related to the length of time the rock is exposed by the tide, which leads to very marked horizontal bands of zonation on most rocky coasts. Supralittoral and littoral fringe zones on the extreme upper shore are lichen dominated. The main eulittoral zone is characterised by barnacles, mussels or furoid algae, the infralittoral by kelps and the circalittoral by animals.	Much less obvious than on rocky coasts, but with a zone of drying on the upper shore and a more water-logged/saturated zone on the lower shore. With increasingly finer sediments the saturated zone extends further up the shore. Very sheltered areas often support saltmarsh vegetation at extreme high water level. Shallow subtidal sediments reflect a high degree of wave disturbance and high temperature/salinity fluctuations, with increasingly more stable conditions with depth.
<i>Exposure to wave action</i>	Marked differences result due to different wave exposures. Exposed shores are usually animal (mussel and barnacle) dominated, whilst sheltered shores are furoid algal dominated. Such differences can occur over only 10's of metres at certain sites, such as opposite sides of a headland. In the subtidal a similar pattern is exhibited, but is increasingly more masked by tidal-current influence with depth.	Principally expressed by the resultant grade of sediment, with coarse sands on exposed coasts and fine muds on sheltered coasts. Areas subject to periodic (seasonal) wave action may exhibit sub-climactic communities.
<i>Strength of tidal currents</i>	Strong offshore currents affect many coasts and have a particularly marked influence on circalittoral communities, with lessening effects in shallow water and on the shore (where the influence of wave action predominates). However constricted sections of some inlets, particularly the narrows in sealochs, can have very strong currents which affect both the shallow subtidal and the lower shore zones, significantly increasing species richness.	Contributes, with wave action, to determining sediment grade and consequent community type. In estuaries and sealochs this can lead to coarser sediments than would normally be expected in wave-sheltered areas. The lower shore of some inlets by the main channel can have tide-swept sands and gravels with distinctive species-rich communities.

Salinity	The majority of rocky coasts are subject to full salinity, but within marine inlets are subject to increasing freshwater influence. Variable salinities (in estuaries) lead to species-poor examples of open coast communities whilst the very limited areas of rock in permanently reduced salinities (in lagoons) may support quite distinct communities. Localised freshwater influence often results in the growth of ephemeral green algae on the shore.	Variable and reduced-salinity conditions are typical of sediment shores within inlets, especially estuaries, and play an important role, alongside sediment type, in determining community type. Salinity eventually becomes the more important structuring factor in the upper reaches of estuaries and in lagoons.
Temperature (relates to biogeography)	National differences in water temperature give more species-rich communities in the south and west and more species-poor communities in the north and east.	
Topography	Topography has a marked influence on the variety of communities which may occur. Variations in topography (resulting from a particular rock type) which lead to vertical faces, overhangs, gullies, caves and rockpools all increase habitat and micro-habitat diversity compared with uniform areas of rock.	Variations in the slope of the beach can indicate differing degrees of saturation, whilst drainage channels may be subject to increased freshwater influence or currents. In the subtidal, variation in slope has little influence on community type, although the presence of dunes can affect small-scale community structure. Larger features such as sandbanks usually indicate coarser sediments compared with surrounding sediment plains.
Geology	The rock type is significant in two respects, affecting overall topography (see above) and the surface texture for colonisation. Soft limestones and chalks have a pitted surface which can affect species composition, whilst these types, plus peats and clays, are soft enough to be bored by piddocks and other species.	Not applicable.
Oxygenation	Not generally applicable, as most rocky habitats are subject to full oxygenation. Severe deoxygenation can lead to reduction in species and the presence of bacterial growths.	More sheltered fine sediments tend to become anoxic below the surface, giving a distinct black layer. Severe deoxygenation significantly reduces species richness.
Wave surge	On exposed coasts gullies subject to wave surge have distinct animal-dominated communities. Wave surge on vertical rock tends to give communities typical of more exposed sites (e.g. <i>Alaria esculenta</i> occurring on moderately exposed vertical rock).	Influences sediment grade and result in highly-mobile species-poor habitats.

Scour, turbidity and siltation	Sand scour and sediment in suspension can encourage growth of ephemeral algae and sometimes mussels (<i>Mytilus</i> spp.) and tube-worms (<i>Sabellaria</i> spp.). Siltation in sheltered areas often restricts the growth of algae.	A high degree of scour and turbidity may result in species-poor communities.
Shading	Shaded faces on the shore encourage the growth of species intolerant of desiccation.	Not applicable.
Organic carbon	Not applicable.	Significant in many sediment communities. Organic enrichment can alter community structure and lead to increased numbers of opportunist species e.g. capitellid worms.
Hydrographic regime (residual currents); water quality	The overall hydrographic regime and water quality characteristics of an area play an important role in determining community composition. Key aspects of these factors are discussed above. In addition to these, residual current flow is also very important, as it may affect larval distribution and water quality aspects such as nutrient levels as well as water temperature, salinity and turbidity.	

Table 2 Summary of anthropogenic influences on community structure

Physical disturbance	Physical disturbance by trampling can impact significantly on rocky shore communities. Disturbance of rock communities in the subtidal is generally less marked. Activities such as fisheries for crabs and lobsters tend to result in only limited changes in the balance of species composition within biotopes but may rarely result in significant shifts in community composition. Where dredging (e.g. for scallops) occurs close to rocky habitats, delicate species can be damaged.	Disturbance of sediment types is widespread, particularly through benthic fisheries activities and aggregate extraction; such disturbance can have significant effects on community composition and may, in some cases, result in completely altered biotopes compared with fully natural conditions. Areas subject to prolonged sedimentary disturbance may exhibit highly variable, sub-climactic communities.
Pollution	Severe pollution may reduce species richness (pollution effects are not well studied).	Pollution may reduce species richness, encourage higher densities of opportunist species, e.g. capitellid polychaetes, or alter community structure.

Classification development - approach and methods used

Review of classification systems and literature

Before embarking on the development of the MNCR BioMar classification (Connor *et al.* 1997a, b), a review of existing classification systems was undertaken (Hiscock & Connor 1991). From these, proposals for a classification structure (Connor *et al.* 1995 a, b) were developed that drew upon the best features of the existing systems, whilst avoiding their weaker aspects. There was subsequent wide consultation on the proposed classification structure, including through two European workshops held during the EC-funded BioMar project (Hiscock *ed.* 1995; Connor *ed.* 1997). These workshops helped ensure broad acceptance of the proposed structure and its wide applicability across European seas.

In addition to a review of classification schemes, an extensive review of the literature describing marine habitats was also undertaken. This helped formulate the initial lists of types which might form the basis of the classification. For this the scientific literature was of considerable help for sediment habitats (a traditional area for marine studies) but relatively poor for rocky habitats (which, in the subtidal, attracted attention only relatively recently through use of SCUBA diving techniques). These initial lists of types were then refined on the basis of new dedicated field surveys, data analyses and field trials.

Consultation and testing

Phases of external consultation and testing of the classification system have been essential to ensure the classification is as robust and usable as possible.

The advice of external consultees has been important in two key areas:

- Marine scientists have contributed expertise in their understanding of the marine environment and its communities, both from a generic perspective and with specific knowledge of communities at particular sites around the country. Of particular importance has been advice on the relationships of environmental factors to community structure and the spatial and temporal dynamics of the marine environment.
- Environmental and conservation managers and end-users have helped define their end needs for the classification system. This has been reflected both in terms of the overall structure of the classification, such as the orientation of biotope complexes to mapping and sensitivity needs, the type of information given in the description of each classification type, and the demands of field application.

Field surveys and other data acquisition

The MNCR undertook a programme of field surveys throughout Britain between 1987 and 1998, collecting data suitable to develop the classification. In addition, data were acquired from the published literature and through collaboration with a wide variety of academic, government and other organisations. Comparable data were collected in Ireland through the BioMar project between 1992 and 1996. The data comprise information on the nature of each site (such as substratum, wave exposure and height or depth surveyed), the type of sampling undertaken, the site's location and the species present (together with an indication of their abundance) within discrete habitats at the site. In total, data for over 16,000 sites comprising more than 36,000 habitat records from around Britain and Ireland have been collated and entered onto the MNCR database. The programme, survey methods and database are described in Hiscock *ed.* (1996). The database includes a module which holds definitions of each classification type, linked to a national dictionary of marine species and to the field survey data. The field survey data have been made widely accessible via the web-based MERMAID application (www.jncc.gov.uk/mermaid) and via the National Biodiversity Network (www.searchnbn.org) from an MS Access-based 'relational' database.

Data analysis

For the 1997 classification, data analyses using the TWINSpan and DECORANA clustering and ordination techniques were employed to help define the types. The analytical processes adopted are described in Mills (1994).

The 1997 version was revised and refined to develop the present version. Extensive re-analyses of the data were carried out using the analytical techniques available in PRIMER (Clarke & Warwick, 2001). The data were initially divided into the five broad habitat types shown in the primary habitat matrix in Table 3 (Littoral Rock, Littoral Sediment, Infralittoral Rock, Circalittoral Rock and Sublittoral Sediment). Due to the large size of the datasets within each broad habitat, some further *a priori* divisions of the data within broad habitats were necessary before analysis was possible. Additional analyses were carried out on data from “borderline” habitats to ensure these *a priori* splits did not force artificial divisions into the classification where this was not supported by differences in the survey data. Analysis within each broad habitat was led by a specialist for that habitat type. The following is a description of the analyses carried out within each broad habitat:

Littoral rock


As the biotopes defined in version 97.06 (Connor *et al.*, 1997 a, b) were generally considered satisfactory, analysis focused on clarifying the boundaries between closely related types and confirming the validity of certain less-well defined types. This included attention to the inter-relationship of fucoid-dominated types regarding the bedrock/boulder/mixed substrata and fully marine/variable salinity transitions and examination of the various red algal-dominated types. Additionally new data from intertidal caves enabled substantial development of the classification here. On the basis of these analyses, some restructuring at biotope complex level was necessary.

Littoral sediment

Due to the size of the Littoral Sediment dataset (>4000 records), some *a priori* division was necessary to provide datasets that could be managed within PRIMER (Clarke & Warwick, 2001). Data were divided based on the sediment type categories at habitat complex level in the 97.06 classification (Connor *et al.*, 1997a, b): gravels and sands, muddy sands, sandy muds, muds and mixed sediments. Semi-quantitative epifaunal data were considered to be of less value than quantitative infaunal data for the purposes of the analysis and were thus excluded. Epifaunal data were however used to define types where a significant proportion of species would be sampled in epibiota sampling techniques, and/or where few infaunal samples were available, e.g. for mussel beds.

Cluster analysis was carried out based on species matrices listing individual counts per m² in each sample, using the PRIMER software package (Clarke & Warwick, 2001). The data were divided into small clusters of biologically similar records, based on the resulting dendrograms. Comparative tables were produced to compare the species data and physical data between each of the small clusters. Where there were no notable differences between the physical and biological characteristics of the small clusters, they were amalgamated into larger groups which would form the preliminary basis for biotopes and sub-biotopes. Where similar biological and physical profiles appeared from clusters derived from different datasets, those data were joined and re-analysed. In particular, there was some overlap between the ‘gravels and sands’ and the ‘muddy sands’, and between the ‘muddy sands’ and ‘mud’ datasets. This re-analysis was carried out to ensure that the *a priori* divisions of the data did not artificially force divisions of otherwise coherent clusters. The resulting preliminary biotope and sub-biotope groups of records were then checked to ensure cohesion of both the environmental and species data. Individual records which differed significantly from the average profile for the group (in terms of biology or physical habitat characteristics) were removed, resulting in a group of records which formed the basis of the biotope descriptions (core biotope records). The physical and biological profiles from the core biotope records were then used to group biotopes of similar character into biotope complexes, and these in turn were assigned to habitat complexes and broad habitats. Note that, in addition to the habitat complexes defined on sediment character, two additional categories were created based on epifaunal characteristics (littoral sediments dominated by macrophytes, and littoral biogenic reefs).

Figure 2. Overview of the data analysis process carried out during classification development (focussing on Littoral Sediments)



Infralittoral rock

As the biotopes defined in version 97.06 were generally considered satisfactory, analysis focused on clarifying the boundaries between closely related types and confirming the validity of certain less-well defined types. This included particular attention to the tide-swept kelp types and the inter-relationship of highly grazed and poorly grazed kelp habitats. On the basis of these analyses, some restructuring at biotope complex level was necessary. Attention was also paid to the vertical rock section of the infralittoral rock classification, and examining how these additional biotopes could be fitted into the existing biotope complexes, reflecting the subtle differences in their biological character.

Circalittoral rock

Due to the complexities of this part of the classification, especially the more subtle differences between types on the open coast, a full re-analysis of the data were undertaken. The large size of the circalittoral rock dataset meant that some *a priori* division was necessary to provide datasets that could be managed within PRIMER (Clarke & Warwick, 2001). Data were divided on the basis of three previously determined energy levels; high, moderate and low energy. Cluster analysis was carried out using epifaunal species matrices exported from the AREV database, using the PRIMER software package (Clarke & Warwick, 2001). The data were divided into small clusters of biologically similar records, based on the resulting dendrograms. Comparative tables were produced to compare the species data and physical data between each of the small clusters. Where there were no notable differences between the physical and biological characteristics of the small clusters, they were amalgamated into larger groups which would form the preliminary basis for biotopes and sub-biotopes. Where similar biological and physical profiles appeared from clusters derived from different datasets, those data were joined and re-analysed. This re-analysis was carried out to ensure that the *a priori* divisions of the data did not artificially force divisions of otherwise coherent clusters. The resulting preliminary biotope and sub-biotope groups of records were then checked to ensure cohesion of both the environmental and species data. Individual records which differed significantly from the average profile for the group (in terms of biology or physical habitat characteristics) were removed, resulting in a group of records which formed the basis of the biotope descriptions (core biotope records). The physical and biological profiles from the core biotope records were then used to group biotopes of similar character into biotope complexes, and these in turn were assigned to habitat complexes and broad habitats. As in the infralittoral rock section, further analysis was also carried out on the vertical rock section of the circalittoral rock classification.

Sublittoral sediment

A full re-analysis of the existing data on the MNCR database in addition to data supplied by the sublittoral specialist was carried out (approximately 10,000 records in total). This followed a similar approach to that described for littoral sediment and as outlined in Figure 2. Data were split according to sediment type, data type (infaunal or epibiota) and sampling technique (where appropriate). Poor quality data was also removed prior to analysis for later manual assessment. Cluster analysis was undertaken using either PRIMER (as described for the littoral sediments) or TWINSpan (following the guidelines in Mills, 1994). Clusters of biologically similar records were produced and assessed using comparative tables. Clusters with poor species definition or highly variable physical characteristics were further sub-divided until more homogenous groups were derived. Where similar biological and physical profiles appeared from clusters derived from different main habitat datasets those data were combined and re-analysed using the same clustering methods as described above in order ensure that the *a priori* divisions of the data did not bias the results of the analysis.

Where similar biological and physical profiles were found in clusters from datasets of differing sampling method or those with different types of data (e.g. epibiota or infauna) the groups were re-analysed where possible at a lower level of resolution (either presence-absence or on the MNCR SACFOR scale) using PRIMER or TWINSpan such that the differences in data type were reduced. As for the littoral sediments the resulting groups were then checked for cohesion with regard the physical and biological data, and individual records assigned to the groups were checked against the profiles of the groups as a whole and re-assigned if necessary.

The physical and biological profiles from the core records for each type were then used to group types of similar character into the broader biotope complexes and these in turn were assigned to one of the six main habitats for sublittoral sediment, derived from the EUNIS classification. The relationship between the sublittoral sediment biotopes is shown for separate depth bands in a series of habitat matrices, available to download as images from the classification website.

Structure of the classification

A framework for the classification (EUNIS levels 2 and 3)

Whilst the classification has been developed for nature conservation purposes and hence needed to be biologically driven, the dynamic nature of certain populations of species, and sometimes whole communities, meant it was essential to identify the habitat within which the community (of potentially varying composition) occurs to ensure types defined would be robust over time. Full use is also made of the habitat attributes to provide a structure to the classification which is both logical and easy to use. In this way much more significant use of habitat characteristics is made than for many terrestrial classifications, where vegetation alone is often the prime determinant of the classification's structure. The classification is presented in such a way as to allow access via either the habitat attributes through a series of habitat matrices or the biological community in a hierarchical classification of biotopes and higher types.

Each of the environmental gradients outlined in Table 1 can be considered to form an axis within a multi-dimensional matrix. Each community develops according to a suite of environmental conditions (and biological influences) which lie within such a multi-dimensional matrix, reflecting varying biological character according to its position along each particular gradient. Although the degree of importance of each habitat attribute varies for differing communities, the first two, namely substratum and the vertical gradient or zonation, appear to play a highly significant role in all communities. They are also the most easily and reliably recorded attributes in the field and are readily mapped. These factors combine to make the attributes of substratum and zonation the most appropriate for structuring the upper end of the classification.

The primary habitat matrix of substrata versus zonation (Table 3) illustrates the framework adopted for the classification. It represents EUNIS levels 2 and 3 in the hierarchical classification and has been developed to reflect the most significant changes in biology at a scale appropriate to an internationally applicable classification. Table 4 outlines the rationale behind the divisions adopted for these two levels in the classification.

Table 3 Framework for the habitat classification - the primary matrix (EUNIS levels 2 and 3). Letters in [] refer to codes. * indicates where various codes are inserted at a lower level in the hierarchy.

SUBSTRATUM		ROCK				SEDIMENT					
		High energy rock [H*R]	Moderate energy rock [M*R]	Low energy rock [L*R]	Features on rock [F*R]	Coarse sediment [CS]	Sand [Sa]	Mud [Mu]	Mixed sediment [Mx]	Macrophyte-dominated sediment [Mp]	Biogenic reefs [BR]
ZONE		(wave exposed or very tide-swept)	(moderately wave-exposed or tide-swept)	(wave sheltered and weak tidal currents)	(rockpools, caves)	Mobile cobble & pebble, gravel, coarse sand	Clean sands & non-cohesive muddy sands	Cohesive sandy muds & muds	Heterogeneous mixtures of gravel, sand & mud		
LITTORAL	LITTORAL [L] (splash zone, strandline & intertidal)	High energy littoral rock [HLR]	Moderate energy littoral rock [MLR]	Low energy littoral rock [LLR]	Features on littoral rock [FLR]	Littoral coarse sediment [LCS]	Littoral sand [LSa]	Littoral mud [LMu]	Littoral mixed sediment [LMx]	Littoral macrophyte-dominated sediment [LMp]	Littoral biogenic reefs [LBR]
	INFRA-LITTORAL [I] (shallow subtidal)	High energy infralittoral rock [HIR]	Moderate energy infralittoral rock [MIR]	Low energy infralittoral rock [LIR]	Features on infralittoral rock [FIR]	Sublittoral coarse sediment [SCS]	Sublittoral sand [SSa]	Sublittoral mud [SMu]	Sublittoral mixed sediment [SMx]	Sublittoral macrophyte-dominated sediment [SMp]	Sublittoral biogenic reefs [SBR]
CIRCA-LITTORAL [C] (nearshore deeper and offshore subtidal)	High energy circalittoral rock [HCR]	Moderate energy circalittoral rock [MCR]	Low energy circalittoral rock [LCR]	Features on circalittoral rock [FCR]							

Table 4 Rationale behind the main divisions adopted in the primary habitat matrix (EUNIS levels 2 and 3)

<i>Rock, Sediment</i>	A primary distinction is made between communities which develop on hard substrata (epibiota) and those which can develop in soft sediments (infauna). Sediments can support distinctive epibiota as well as infauna. The term rock is used in a broad sense to indicate hard substrata such as bedrock, boulders, stable cobbles, artificial substrata and biogenic substrata. Sediments also include pebbles and cobbles which are essentially mobile (shingle) or may have a small proportion of stones and shells on the surface, supporting epibiota. Where biogenic substrata develop on substantially sediment substrata, they are included in the sediment section of the classification.
<i>Littoral, Sublittoral (Infralittoral, Circalittoral)</i>	These represent the major divisions in a vertical gradient from the terrestrial environment to the edge of the continental shelf (about 200 m depth). The main factors which control the zonation are immersion, thermal stability, light, wave action and salinity. They interact in a complex manner to produce a general zonation pattern, applicable to both rock and sediment habitats throughout Europe and beyond. Table 5 illustrates the inter-relationship of the factors for each zone, and Figure 1 provides a typical schematic profile of this zonation pattern.
<i>High energy rock, Moderate energy rock, Low energy rock</i>	These are defined on an energy gradient, reflecting exposure to wave action or tidal currents, or a combination of both (note, this energy gradient was reflected in the 1997 classification, but expressed as 'exposure'; the resulting confusion with wave exposure has now been removed). This energy gradient is broadly paralleled in sediment habitats, where coarse clean sediments occur in high energy conditions and fine muds occur in low energy conditions. Although the effects of wave action and tidal currents can be significantly different, there are many instances where the increase in tidal current strength in wave-sheltered habitats gives rise to communities similar to those found on more wave-exposed coasts but in reduced tidal currents. For example, increased currents in the infralittoral zone change the kelp <i>Laminaria saccharina</i> communities of very wave-sheltered sites to <i>Laminaria hyperborea</i> communities similar to those on open, more wave-exposed coasts. Very strong tidal currents in the circalittoral appear to override the effect of wave action to a large extent, giving rise to a suite of associated communities of barnacles, cushion sponges and the hydroid <i>Tubularia indivisa</i> which are less obviously affected by wave action. These communities are similar in character to those of surge gullies which are subject to extreme wave action.

***Coarse sediments,
Sands, Muds,
Mixed sediments,
Macrophyte
communities on
sediments,
Biogenic Reefs***

The particular sediment grade, typically derived from the hydrodynamic conditions of the site, strongly influences community structure. The four main divisions adopted here reflect major changes in species character, particularly related to the amount of silt or clay in the sediment. In addition, some sediments support communities of macrophytes (angiosperms and seaweeds) which attach to small stones and shells on the sediment surface, whilst on others biogenic reefs develop in which a particular species aggregates to form a stable surface upon which other species can live. With both macrophyte and biogenic reef communities the underlying sediment may support infaunal communities according to the particular sediment type; however the prominent character of the epibiota communities has led to a preference to group such biotopes under these separate major categories.




Figure 1 Profile of a rocky shore and seabed showing the biological zones (modified from Hiscock *ed.* 1996); heights and depths given are typical values for south-west Britain. In sediment habitats a similar vertical zonation for the main zones is found.

Table 5 Marine biological zones and the factors determining them

<i>Zone</i>	<i>Typical upper boundaries around Britain and Ireland</i>	<i>Immersion</i>	<i>Thermal stability</i>	<i>Light</i>	<i>Salinity</i>	<i>Wave action</i>
<i>Adlittoral</i>		Spray only	Highly variable	Photic	Saline influence	None
<i>Supralittoral</i>	+10 to +6 m	Spray and splash	Highly variable	Photic	Euryhaline	Highly variable
<i>Eulittoral</i>	+7 to +4 m MHWS	Regular immersion and emersion	Highly variable	Photic	Euryhaline	Highly variable
<i>Infralittoral</i>	+1 to 0 m MLWS	Immersed (intermittent spring tide emersion of sublittoral fringe)	Variable - eurythermal	Euphotic	Euryhaline	Variable
<i>Circalittoral</i>	-5 to -20 m	Immersed	Moderately variable - mesothermal	Mesophotic (sparse algae, algal crusts)	Mesohaline / Stenohaline	Moderately variable
<i>Circalittoral offshore</i>	-40 to -80 m	Immersed	Stable - stenothermal	Aphotic	Stenohaline	Stable
<i>Bathyal</i>	-200 m	Immersed	Very stable - stenothermal	Aphotic	Stenohaline	Stable

Development of a hierarchical classification

It was considered essential to develop a hierarchical classification structure in which broader, higher types in the classification could be more finely divided to support more detailed use. The development of the hierarchy comes from both a top-down and a bottom-up approach:

Top-down classification

Taking the marine environment as a whole, it can be sub-divided into a series of broad habitat categories, based largely on their physical character as described here. At the very broadest level, differentiation can be made between rock and sediment habitats, and between those on the shore (intertidal) and those in the subtidal or deep ocean. These high-level divisions can be further subdivided on the basis of different types of sediment (e.g. gravel, mud), different degrees of wave exposure on rocky coasts (exposed, sheltered) and varying depth bands below the low water mark (e.g. shallow water where light penetrates, deeper water with little light). Such broad-scale differences in habitat character are readily understood by non-specialists and provide classification types that are easily mapped; however, they also have ecological relevance as they reflect major changes in habitat character upon which species depend (see above).

The top-level types depicted in the primary habitat matrix (Table 3) show levels 2 and 3 in the hierarchical classification. It is important to note that these top-level categories were developed after consideration of how best to classify biological data at the lower end of the classification.

Bottom-up classification

Field survey, whether on the shore or in the subtidal, reveals that different places support different communities of species. The precise combination of species and their relative abundance varies from place to place and is dependent both on environmental characteristics and upon interactions between

species. Visits to different sites that have similar environmental characteristics, such as sediment type and depth, show certain levels of similarity in their species composition. Multivariate analysis of the data from field surveys (e.g. grabs, diver observations) groups these data into clusters that have similar character – this forms the basis of defining the types at the lower end of the classification (levels 5 and 6). These can themselves be grouped into higher types with similar character (level 4), thus forming the basis for the bottom-up approach to development of the classification based on real field sample data.

The two approaches have been merged together into a single hierarchy, thus catering for broad-scale application in management and mapping and fine-scale application for detailed survey, monitoring and scientific study. The levels can be differentiated in relation to their degree of biological distinctiveness, to the ability to discriminate types by various methods of remote and *in situ* sampling, to the ease of recognition by workers with differing skill levels and to the end use of the classification for conservation management at various scales.

Six levels in the hierarchy have thus been developed, equating directly to the levels in the EUNIS classification:

Level 1 Environment (marine) – A single category is defined within EUNIS to distinguish the marine environment from terrestrial and freshwater habitats.

Level 2 Broad habitats - These are extremely broad divisions of national and international application for which EC Habitats Directive Annex I habitats (e.g. reefs, mudflats and sandflats not covered by seawater at low tide) are the approximate equivalent.

Level 3 Main habitats - These serve to provide very broad divisions of national and international application which reflect major differences in biological character. They are equivalent to the intertidal Sites of Special Scientific Interest (SSSI) selection units (for designation of shores in the UK) (Joint Nature Conservation Committee 1996) and can be used as national mapping units.

Level 4 Biotope complexes - These are groups of biotopes with similar overall physical and biological character. Where biotopes consistently occur together and are relatively restricted in their extent, such as rocky shores and very near-shore subtidal rocky habitats, they provide better units for mapping than the component biotopes, better units for management and for assessing sensitivity than the individual biotopes. They are relatively easy to identify, either by non-specialists or by coarser methods of survey (such as video or rapid shore surveys), thereby offering opportunities for data collection by a wide range of people and without recourse to specialist species identification skills.

Level 5 Biotopes - These are typically distinguished by their different dominant species or suites of conspicuous species. On rocky substrata, most should be readily recognised by workers with a basic knowledge of marine species, although quantitative sampling will be necessary in many of the sediment types. The vast majority of available biological sample data are attributable to this level (or the sub-biotope level), which is equivalent to the communities defined in terrestrial classifications such as the UK National Vegetation Classification (e.g. Rodwell *ed.* 1995). Intertidal and subtidal sediment biotopes may cover very extensive areas of shore or seabed.

Level 6 Sub-biotopes - These are typically defined on the basis of less obvious differences in species composition (e.g. less conspicuous species), minor geographical and temporal variations, more subtle variations in the habitat or disturbed and polluted variations of a natural biotope. They will often require greater expertise or survey effort to identify.

The primary habitat matrix (Table 3) provides an overview of levels 2 and 3 in the classification. Biotope matrices for each broad habitat, showing the relationship of biotopes and sub-biotopes to key environmental factors, can be downloaded as images from the classification website. For each broad habitat, a hierarchy structure diagram showing the relationship between units at the higher and lower hierarchical levels has been created in Excel™, and all the types in the classification are listed in hierarchical order in an additional Excel™ spreadsheet, all available to download from the classification website

Distinguishing and defining types

To ensure consistency across the classification in how types are defined, a working definition as to what constitutes a biotope, enabling its distinction from closely-related types, has been developed. The following criteria are applied:

1. The entity can be distinguished on the basis of a consistent difference in species composition based on:
 - different dominant species, some of which (e.g. mussels and kelps) may be structurally important; and
 - the co-occurrence of several species characteristic of the particular habitat conditions (even though some of these may occur more widely in other combinations).

A combination of both the presence and abundance of the most 'obvious' species in a community is used. Sub-biotopes are often defined using less conspicuous species.
2. It occurs in a recognisably different habitat (but acknowledging that distinct communities may develop in the same habitat through change with time). Sub-biotopes are often defined on the basis of more subtle habitat differences. Some highly subtle differences may be critical in determining community structure (e.g. water circulation/exchange patterns in sealoch basins, oxygenation levels in the water column/sediment, sediment structure other than grain size composition). The separate divisions of habitat factors used in field recording are not necessarily be reflected in the end division of types.
3. It is a recognisable entity in the field, i.e. it is not an artefact of data analysis.
4. The assemblage of species recurs under similar habitat conditions in (at least several) widely-separate geographical locations. Associations of species confined to a small geographical area are considered unlikely to represent a recurrent community (unless the habitat is considered unique), but should rather be treated as a variation of a more widely occurring type.
5. As a working guide the biotope extends over an area at least 5 m x 5 m, but can also cover many square kilometres, such as for extensive offshore sediment plains. For minor habitats, such as rockpools and overhangs on the shore, this 'minimum size' can be split into several discrete patches at a site. Small features, such as crevices in rock or the biota on kelp stipes, are described as features of the main biotope rather than biotopes in their own right. Some entities, by virtue of their extent around the coast, may warrant description despite showing only minor differences in species composition; such types are often treated as sub-biotopes.
6. It is a single entity in the field, although there may be some spatial variation or patchiness from one square metre to the next. Therefore each area of shore or seabed should correlate to only one biotope defined in classification (a 1:1 relationship of field units to classification units). Whenever possible, the surface species characteristics of sediment habitats (their epibiota) are described in association with the sediment infauna as a single entity, rather than treated as separate communities. Note however that the nature of available data has severely restricted the clear association of these two aspects in the classification as they are typically derived from differing survey techniques. Thus in the present classification there remain units defined primarily on the basis of their epibiota or their infauna but which, given further research, will be shown to be the same biotope. Epibiota-derived biotopes may also 'overlay' a number of infaunal biotopes, which are differentiated by more subtle environmental differences, and thus need to be referred to a higher unit in the classification.

The following considerations are also taken into account in deciding whether to establish a biotope:

- There is a need to recognise that it is commonplace to have no distinct boundary between two different 'types', but a gradual transition, such that distinction of types is somewhat arbitrary at particular reference points or nodes along a continuum. Additionally, some communities may be largely transitional (in a temporal sense) in nature and whilst recognisable in the field represent a stage between two or more 'stable' biotopes. In some areas, e.g. due to periodic

disturbance, a community may be held in a transitional or sub-climactic state for prolonged periods and certain habitats may be so variable that the position of a biotope along a gradient cannot be accurately defined. These factors are of critical importance when assessing typicality of a site to a particular type or its quality or conservation importance.

- Where different associations are shown to occur within the same habitat, they may be spatial or temporal mosaics caused by factors such as grazing, disturbance or chance recruitment. These should be linked together in the classification as, for conservation purposes, it is important to manage or protect the habitat in which several communities may occur over time.
- To produce a practicable working classification it has been necessary at times to be general rather than specific in splitting different types, so that an excessively and unnecessarily complex classification is not developed (bearing in mind the end units that are necessary for practical use).
- Separation of communities can be related to conservation value - does the type add variety (of habitat or species) to a particular stretch of coast. This relates to natural habitats and excludes artificial, polluted or disturbed habitats which should not be considered of high conservation value although they may support distinct communities.

How to use the classification

Habitat matrices

The primary habitat matrix (Table 3) provides a general framework for the classification and shows the level 2 and 3 types. In addition, more detailed matrices have been created for each of the broad habitats, showing the distribution of individual biotopes and sub-biotopes (levels 5 and 6) in relation to key habitat factors. The format of these matrices has changed significantly from previous versions of the classification to better depict the relationship of biotopes to habitat factors. For the rocky habitats, biotopes are shown in relation to energy levels, whereas for sediment habitats, biotopes are shown in relation to sediment type using a modified Folk triangle approach (Folk 1954).

The matrices aim to provide a rapid indication of the range of biotopes that could occur under particular habitat conditions, e.g. moderate energy infralittoral rock or intertidal sandflats. They can be used to indicate which closely related biotopes should be considered before determining to which type a sample record should be assigned.

Presentation of the biotopes and sub-biotopes within these matrices has a number of benefits:

- It helps to display the relationship of a biotope to other closely related types and to clarify the main habitat parameters which contribute to its structure. These relationships are less clear in a more conventional listing of types (e.g. the hierarchical listing).
- It enables the identification of dissimilar communities within apparently similar physical environments. Here, although there may be subtle physical factors which drive such differences in biological composition, other factors such as seasonal change, chance recruitment, grazing pressures or pollution effects may account for the differences and allow such communities to be linked within the classification.
- It also facilitates the undertaking of new ecological survey in a more structured manner, by enabling the full range of habitats in an area to be identified and sampled.

The matrices for littoral rock, littoral sediment, infralittoral rock, circalittoral rock and sublittoral sediment are available to download as images from the classification website.

Hierarchy structure diagrams

Hierarchy structure diagrams have been created for each of the broad habitats, providing an overview of all the lower level units within each broad habitat type. These diagrams are available to download as Excel™ spreadsheets from the classification website. They have also been included at the end of the pdf documents containing the biotope descriptions available to download for each section of the classification.

Hierarchical list of types

A full hierarchical list of types is available to download as an Excel™ spreadsheet from the classification website. The types are presented in hierarchical order, to help bring together those types which are most similar to each other in character.

Layout of descriptions for each type

Descriptions for each unit in the classification, from broad habitats to sub-biotopes, are laid out as follows:

Code

A unique letter code, reflecting the level of the described type within the classification hierarchy.

Title

The title gives the key biological and physical features of the type, with emphasis on the features which help to distinguish it from closely related types of the same level in the hierarchy. The habitat

part of the title usually includes the zone, substratum and another key habitat factor. To avoid becoming overly clumsy the titles do **not** cover all habitat characteristics or characterising species, and common names are not given (although they are given in the text description).

NOTE: It is very important to refer to the full description and to the habitat matrices to determine the full nature of the type and not to rely on the title alone.

Habitat characteristics

The typical habitat characteristics of the type for salinity, wave exposure, tidal currents, substratum, zone, height or depth band and, where appropriate, other factors critical to that particular type. The range given for each factor tends to be broader for higher types and more tightly defined for lower types. When assigning samples to types, it should be noted that in some cases the type may occur outside the range given (see profiles given in the comparative tables which show that a small proportion of records may occur outside the typical range for the type), though care should be taken to ensure that another type has not been described to cover the example being considered. All heights and depths are corrected to chart datum.

Previous code

Codes used in versions 6.95, 96.7, and 97.06 (Connor *et al.* 1995 a, 1996, 1997 a, b) are given where different to the current code. Where communities from previous versions have been combined or split, previous codes are shown as far as possible. Some communities in the revised classification are newly defined and may not relate directly to types in the previous classification. Lookup tables which help to translate 2004 to 1997 codes (and vice versa) are available to download in the form of an Excel™ spreadsheet from the classification website.

Description

An account of the general nature of the habitat and community characteristics, and its micro-habitat features (e.g. crevices, under-boulders, kelp stipes) if present.

Situation

Describes the general situation on the shore or in the sublittoral, in relation to other types (i.e. along gradients of substratum, zonation, wave exposure, tidal currents, salinity etc.).

Temporal variation

This section outlines the known natural temporal dynamics of the type described, such as seasonal changes in community structure or physical environment. In general, much more information is needed for this section. In some cases separate types may have been defined because there is a lack of knowledge that the communities are temporal variations within a single habitat type.

Similar types

Attention is drawn to similar types which should be considered before assigning a field record to a particular biotope. The main similarities and principal distinguishing features are described for each similar biotope.

Characterising species

A list of those species which contribute most to the overall similarity between core records assigned to the type, i.e. characterise the type, with associated information on their frequency of occurrence, their individual contribution to the similarity within the core data set of records, and the typical abundance at which they occur.

For each type, characterising species have been determined using the SIMPER routine in PRIMER (Clarke & Warwick, 2001). For a given set of records (in this case, core records of each type), SIMPER indicates and ranks the individual contribution of each species to the overall similarity within the data set. Both the frequency of occurrence of each species within the dataset and their abundance (using the SACFOR abundance scale (p.46) for epifaunal data and numeric counts for infaunal data)

are taken into account during this process. Species that contribute more than 1% to the overall similarity of the records within the data set are defined as 'characterising species', and listed in a characterising species table. Those that contribute less than 1% are not listed. Species which qualify according to the SIMPER routine, but are Present or Rare on the MNCR SACFOR scale and present in fewer than 20% of the records, are occasionally excluded from the characterising species table.

Care has been taken to mention each of the characterising species in the descriptions for each type. Sometimes additional species are mentioned that are particularly indicative (faithful) of that type or characteristic of a biogeographic region, but which have not qualified as 'characterising species' according to the SIMPER routine.

The **% contribution to similarity** column of the table shows the contribution of each characterising species to the similarity within the type, i.e. the higher the contribution, the higher the importance of the species. The number of species in the table reflects the species diversity within each type. In types with a high species richness, a large number of species each contribute with a relatively low amount to the similarity within the group. If a type has low diversity, then a small number of species contribute with relatively large amounts to the overall similarity and hence fewer species are listed in the table. In a few cases, a long species list indicates low overall similarity of records within the type.

The **% frequency of occurrence** column of the table shows the occurrence of a species within a certain biotope. The symbols represent percentage occurrence in the samples as follows:

- Occurs in 81-100% of the records for the type
- Occurs in 61-80% of the records for the type
- Occurs in 41-60% of the records for the type
- Occurs in 21-40% of the records for the type

The **typical abundance** column of the table shows the mean SACFOR abundance for each characterising species within the samples where it is present. Quantitative infaunal counts have been converted to the SACFOR scale for compatibility of data presentation. For types where the core records are exclusively quantitative infaunal records (e.g. most of the littoral sediment types), an additional column is included in the characterising species table, showing mean counts per m² for each species within the core data set.

Comparative tables

Comparative tables of physical (habitat) and biological (species) data have been generated in the form of Excel™ spreadsheets to enable a comparative overview over similar biotopes or other classification units. The spreadsheets are available to download on the classification website.

Comparative tables of habitat data

The habitat (physical) comparative tables are generated from the MNCR database, using only the core data sets for each type (see data analysis section). Each column represents a separate type.

The left column of the table lists each of the habitat factors and their categories, as per the field data. For each type, the percentage of records containing data for each category is shown both in figures and as a bar. For example a table may indicate that 85% of the records for a type were recorded as full salinity and the remaining 15% as variable salinity.

The percentage given reflects only the available data for that factor. Consequently, where the data for a factor are lacking, particularly if that is a high proportion of full data set for that data, the resulting percentage distribution across the categories may not fully reflect the character of that type.

Where more than one category can be recorded in the field data, e.g. zone and depth band, the figures within a habitat factor may exceed 100%.

Comparative tables of species data

The species comparative tables are generated from the MNCR database, using only the core data sets for each type (see data analysis section, p.17 ff). Each column represents a separate type.

The left column of the table lists, in taxonomic order, the species which occur in at least one of the types shown in that table. To ensure clarity, the tables are compiled with a minimum percentage cut-off of 20%: that is, only species present in more than 20% of the records for a particular type are displayed. The percentage of records for each type containing each species is shown both in figures and as a bar.

For types where the core records are semi-quantitative samples (e.g. epifaunal samples on rocky shores), a letter alongside these bars indicates the median SACFOR abundance of that species from the records that it occurred in. For example, C 50 next to *Patella vulgata* indicates that *P. vulgata* occurred in 50% of the records for that particular type and that it had a median abundance of “common” where it occurred.

For types where the core records consist of quantitative samples (e.g. infaunal core samples on intertidal sediments), the comparative tables display the mean number of individuals per m².

Species nomenclature

All species names are given according to Howson & Picton (1997), excepting for angiosperms, which follow Stace (1991), and lichens, which follow Purvis *et al.* (1992). Guiry & Dhonncha (2002) provides a later checklist for algae and additional useful information; the present publication and database does not yet follow this revised checklist.

Understanding the codes

A letter coding system has been adopted in preference to a fully numerical coding system or an alphanumeric system (as used in the NVC and EUNIS systems). This has a number of advantages. It enables the construction of intuitive codes which can readily be related to their respective types without recourse to the full type title. Furthermore, it enables changes to the order in which the types are presented without the need to change a numerically sequenced code. This was particularly useful in the early development phase of the classification, but has continued to be of use during subsequent revisions of the classification.

Construction of codes follows a few simple rules, which achieve consistency throughout the classification whilst aiming to keep the resultant codes relatively short and intuitive. Familiarity with the rules for code construction and with the types themselves, by those working regularly with the classification, results in rapid use of codes as a short-hand means of referring to the types defined.

Codes are defined for each level in the classification. Within a level, they comprise one or several elements. They are based on the following rules:

1. Broad habitat and main habitat codes are based on habitat factors or gross biological features (e.g. macrophytes and biogenic reefs).
2. Biotope complex, biotope and sub-biotope codes are based wherever possible upon the most characteristic taxa (which preferably also dominate spatially/numerically) (preferably no more than two per biotope complex, biotope or sub-biotope).
3. Where the biological composition is too complex to derive a simple code, features of the habitat are used (e.g. VS for variable salinity).
4. Codes for habitat factors, higher taxa and descriptive community features (e.g. park, crustose) are derived from a standard lexicon (Table 6). A full list of codes used is contained in the hierarchical list which can be downloaded from the classification website.
5. Codes for names of genera are derived using the first three letters of a genus or higher taxon name (e.g. Ala for *Alaria*, Chr for *Chrysophyceae*). Codes for species names are derived using the first letter of the genus and the first three letters of the specific name (e.g. Ldig for *Laminaria digitata*).
6. Within the code each new element of the code starts with a capital letter.

7. As far as practical the code elements are unique, but some duplication is adopted in the interests of keeping codes short. The code for any given type (i.e. for the level defined, regardless of whether it is stringed with higher codes – see below) is always unique.
8. All the biotope/sub-biotope codes are unique, so users familiar with the classification can refer to individual biotopes using only the codes for these levels in the hierarchy.
9. The full codes are compiled using the code for each level in the hierarchy, separated from the next level by a full stop, starting with the broad habitat (level 2), followed by the main habitat, biotope complex and so on. For example LS.LSA.MoSa.AmSco.Eur:

2	broad habitat	littoral sediment	LS
3	main habitat	littoral sand	LSA
4	biotope complex	mobile sand	MoSa
5	Biotope	Amphipods and <i>Scolelepis</i> spp.	AmSco
6	Sub-biotope	<i>Eurydice</i> sub-biotope	Eur

NOTE: to avoid confusion, others using the classification should not erect similar codes for types not currently described in the national classification.

Table 6 Lexicon of codes (excluding those at genus and species level)

Code	Meaning	Type	Level	97.06 code	Comments
Aalb	<i>Abra alba</i>	Genus/species	4, 5, 6	Abr	
Aasp	<i>Ascidiella aspersa</i>	Genus/species	4, 5, 6	Aasp	
Abr	<i>Abra</i>	Genus/species	4, 5, 6		
Abra	<i>Amphiura brachiata</i>	Genus/species	4, 5, 6		
Achi	<i>Amphiura chiajei</i>	Genus/species	4, 5, 6	Achi	
Act	<i>Actinothoe</i>	Genus/species	4, 5, 6		
Adia	<i>Alcyonidium diaphanum</i>	Genus/species	4, 5, 6	Adia	
Adig	<i>Alcyonium digitatum</i>	Genus/species	4, 5, 6	Alc	
Afal	<i>Ampharete falcata</i>	Genus/species	4, 5, 6	Amp	
Afil	<i>Amphiura filiformis</i>	Genus/species	4, 5, 6	Afil	
Aglo	<i>Alcyonium glomeratum</i>	Genus/species	4, 5, 6		
Ahn	<i>Ahnfeltia</i>	Genus/species	4, 5, 6	Ahn	
Airr	<i>Astropecten irregularis</i>	Genus/species	4, 5, 6		
Al	Algae/algal	Taxon group	4, 5, 6	Al	
Ala	<i>Alaria</i>	Genus/species	4, 5, 6	Ala	
Am	Amphipods	Taxon group	4, 5, 6	A	
Amen	<i>Ascidia mentula</i>	Genus/species	4, 5, 6	Amen	
Amp	<i>Ampelisca</i>	Genus/species	4, 5, 6		
Amy	<i>Amythasides</i>	Genus/species	4, 5, 6		
An	Anemones	Taxon group	4, 5, 6	An	
Ang	Angiosperms	Taxon group	4, 5, 6	Ang	
Anit	<i>Abra nitida</i>	Genus/species	4, 5, 6		
Ant	<i>Antedon</i>	Genus/species	4, 5, 6	Ant	

Aope	<i>Aequipecten opercularis</i>	Genus/species	4, 5, 6		
Aph	<i>Aphelochaeta</i>	Genus/species	4, 5, 6	Aph	
Apri	<i>Abra prismatica</i>	Genus/species	4, 5, 6		
Aps	<i>Apseudes</i>	Genus/species	4, 5, 6		
Are	<i>Arenicola</i>	Genus/species	4, 5, 6	Are	
As	Ascidians	Taxon group	4, 5, 6	As	
Asc	<i>Ascophyllum</i>	Genus/species	4, 5, 6	Asc	
Ascmac	<i>Ascophyllum nodosum ecad mackaii</i>	Genus/species	4, 5, 6	Asc*mac	
Asqu	<i>Amphipholis squamata</i>	Genus/species	4, 5, 6		
Aten	<i>Angulus tenuis</i>	Genus/species	4, 5, 6		
Aud	<i>Audouinella</i>	Genus/species	4, 5, 6		
Axi	Axinellid sponges	Taxon group	4, 5, 6	Axi	
B	Barnacles	Taxon group	4, 5, 6	B	
B	Biogenic [reefs]	Community feature	2, 3		
Bal	<i>Balanus</i>	Genus/species	4, 5, 6	Bal	
Bar	Barren	Community feature	4, 5, 6	Bar	
Bat	<i>Bathyporeia</i>	Genus/species	4, 5, 6	Bat	
Beg	<i>Beggiatoa</i>	Genus/species	4, 5, 6	Beg	
Bif	<i>Bifurcaria</i>	Genus/species	4, 5, 6	Bif	
Blan	<i>Branchiostoma lanceolatum</i>	Genus/species	4, 5, 6	Bra	
Bli	<i>Blidingia</i>	Genus/species	4, 5, 6	Bli	
Blyr	<i>Brissopsis lyrifera</i>	Genus/species	4, 5, 6	Bri	
Bo	Boulders	Habitat factor	4, 5, 6	Bo	
Br	Brachiopods	Taxon group	4, 5, 6	Br	
Bri	Brittlestars	Taxon group	4, 5, 6	Bri	
Bug	<i>Bugula</i>	Genus/species	4,5,6	Bug	
By	Bryozoans	Taxon group	4, 5, 6	By	
C	Circalittoral	Habitat factor	2, 3	C	
C	Coarse [sediment]	Habitat factor	2, 3		
Cap	<i>Capitella</i>	Genus/species	4, 5, 6	Cap	
Cape	Cape-form (kelp)	Community feature	4, 5, 6		
Car	<i>Caryophyllia</i>	Genus/species	4, 5, 6	Car	
Care	<i>Corophium arenarium</i>	Genus/species	4, 5, 6	Cor	
Cb	Cobble	Habitat factor	4, 5, 6		
CC	Crustose coralline algae	Taxon group	4, 5, 6	CC	
Ccas	<i>Cordylophora caspia</i>	Genus/species	4, 5, 6	Cor	
Ccor	<i>Clathrina coriacea</i>	Genus/species	4, 5, 6	Cla	
Cer	<i>Cerastoderma</i>	Genus/species	4, 5, 6	Cer	
Cha	<i>Chara</i>	Genus/species	4, 5, 6		
Cho	<i>Chorda</i>	Genus/species	4, 5, 6	Cho	
Chr	Chrysophyceae	Taxon group	4, 5, 6	Chr	

Cht	<i>Chthamalus</i>	Genus/species	4, 5, 6	Cht	
Cio	<i>Ciona</i>	Genus/species	4, 5, 6	Cio	
Cir	Cirratulid polychaetes	Taxon group	4, 5, 6		
Cla	<i>Cladophora rupestris</i>	Genus/species	4, 5, 6		
Cllo	<i>Cerianthus lloydii</i>	Genus/species	4, 5, 6		
Co	Colonial [ascidians]	Community feature	4, 5, 6		
Cod	<i>Codium</i>	Genus/species	4, 5, 6	Cod	
Coff	<i>Corallina officinalis</i>	Genus/species	4, 5, 6	Coff & Cor	
Con	<i>Conopeum</i>	Genus/species	4, 5, 6	Con	
Cor	Corallinaceae/coralline	Taxon group	4, 5, 6	Cor	
Cr	Crusts/crustose	Community feature	4, 5, 6	C	
Cre	<i>Crepidula</i>	Genus/species	4, 5, 6	Cre	
Cri	Crisiid bryozoans	Taxon group	4, 5, 6	Cri	
Crl	Coral (reefs e.g. <i>Lophelia</i>)	Taxon group	4, 5, 6		
CrSp	Crustose sponges	Taxon group	4, 5, 6	SC	
Cset	<i>Chaetozone setosa</i>	Genus/species	4, 5, 6		
Cu	Cushion [sponges]	Community feature	4, 5, 6	CuS	
Cum	Cumaceans	Taxon group	4, 5, 6		
Cup	Cup corals (Scleractinia)	Taxon group	4, 5, 6	Cup	
Cv	Caves	Habitat factor	4, 5, 6	Cv	
Cvar	<i>Chlamys varia</i>	Genus/species	4, 5, 6	Cvar	
Cvir	<i>Corynactis viridis</i>	Genus/species	4, 5, 6	Cor	
Cvol	<i>Corophium volutator</i>	Genus/species	4, 5, 6	Cor	
Cys	<i>Cystoseira</i>	Genus/species	4, 5, 6	Cys	
Den	<i>Dendrodoa</i>	Genus/species	4, 5, 6	Den	
Des	<i>Desmarestia</i>	Genus/species	4, 5, 6		
Dic	<i>Dictyopteris</i>	Genus/species	4, 5, 6	Dic	
Dp	Deep (circalittoral)	Habitat factor	4, 5, 6		
Dys	<i>Dysidia</i>	Genus/species	4, 5, 6		
Ec	Echinoderms	Taxon group	4, 5, 6		
Ecor	<i>Echinocardium cordatum</i>	Genus/species	4, 5, 6	Ecor	
Edef	<i>Eudorellopsis deformis</i>	Genus/species	4, 5, 6		
Edw	Edwardsia	Genus/species	4, 5, 6	Edw	
Ele	<i>Electra</i>	Genus/species	4, 5, 6	Ele	
Ens	<i>Ensis</i>	Genus/species	4, 5, 6	Ens	
Ent	<i>Enteromorpha</i>	Genus/species	4, 5, 6	Ent	
Eph	Ephemeral (seaweeds)	Community feature	4, 5, 6	Eph	
Epus	<i>Echinocyamus pusillus</i>	Genus/species	4, 5, 6		
Er	Erect [sponges]	Community feature	4, 5, 6	ErS	
Est	Estuarine	Habitat factor	4, 5, 6	Est	

Ete	<i>Eteone</i>	Genus/species	4, 5, 6		
Eud	<i>Eudendrium</i>	Genus/species	4, 5, 6	Eud	
Eun	<i>Eunicella</i>	Genus/species	4, 5, 6	Eun	
Eur	<i>Eurydice</i>	Genus/species	4, 5, 6	Eur	
F	Features (e.g. rockpools, caves)	Habitat factor	2, 3		
F	Fucoids	Taxon group	4, 5, 6	F	
F	Full [salinity] (=marine)	Habitat factor	4, 5, 6	FS	
Fa	Fauna/faunal	Taxon group	4, 5, 6	Fa	
Fab	<i>Fabricia</i>	Genus/species	4, 5, 6	Fab	
Fcer	<i>Fucus ceranoides</i>	Genus/species	4, 5, 6	Fcer	
Fdis	<i>Fucus distichus</i>	Genus/species	4, 5, 6	Fdis	
Ffab	<i>Fabulina fabulina</i>	Genus/species	4, 5, 6		
Fi	Fine [sand or mud]	Habitat factor	4, 5, 6		
Fil	Filamentous (seaweeds)	Community feature	4, 5, 6	Fi	
Flu	<i>Flustra</i>	Genus/species	4, 5, 6	Flu	
Fo	Foliose (seaweeds)	Community feature	4, 5, 6	Fo	
For	Foraminiferans	Taxon group	4, 5, 6	For	
Fou	Fouling	Community feature	4, 5, 6		
Fser	<i>Fucus serratus</i>	Genus/species	4, 5, 6	Fser	In MLR
Fserr	<i>Fucus serratus</i>	Genus/species	4, 5, 6	Fserr	In LLR
Fspi	<i>Fucus spiralis</i>	Genus/species	4, 5, 6	Fspi	
Ft	Forest (kelp)	Community feature	4, 5, 6	Ft	
Fun	<i>Funiculina</i>	Genus/species	4, 5, 6	Fun	
Fur	<i>Furcellaria</i>	Genus/species	4, 5, 6	Fur	
Fves	<i>Fucus vesiculosus</i>	Genus/species	4, 5, 6	Fves	
G	Green seaweeds (Chlorophyceae)	Taxon group	4, 5, 6	G	
G	Gully [surge gully]	Habitat factor	4, 5, 6	G	
Gam	<i>Gammarus</i>	Genus/species	4, 5, 6		
Glap	<i>Glycera lapidum</i>	Genus/species	4, 5, 6		
Gra	<i>Gracilaria</i>	Genus/species	4, 5, 6		
Gv	Gravel/gravelly	Habitat factor	4, 5, 6		
Gz	Grazed (seaweed communities)	Community feature	4, 5, 6	Gz	
H	High energy (very wave/tide exposed)	Habitat factor	2, 3	E	
H	Hydroids	Taxon group	4, 5, 6	H	
Hal	<i>Halidrys</i>	Genus/species	4, 5, 6	Hal	
Hap	Haptophyceae	Taxon group	4, 5, 6		
Har	<i>Hartlaubella</i>	Genus/species	4, 5, 6	Har	
Hbow	<i>Halichondria bowerbanki</i>	Genus/species	4, 5, 6	Hbow	

Hchr	<i>Halcampa chrysanthellum</i>	Genus/species	4, 5, 6	Hal	
Hed	<i>Hediste</i>	Genus/species	4, 5, 6	Hed	
Helo	<i>Hesionura elongata</i>	Genus/species	4, 5, 6		
Het	<i>Heteromastus</i>	Genus/species	4, 5, 6		
Hia	<i>Hiatella</i>	Genus/species	4, 5, 6	Hia	
Hil	<i>Hildenbrandia</i>	Genus/species	4, 5, 6		
Him	<i>Himantalia</i>	Genus/species	4, 5, 6	Him	
Ho	Holothurians	Taxon group	4, 5, 6	Ho	
Hocu	<i>Haliclona oculata</i>	Genus/species	4, 5, 6	Hocu	
Hyd	<i>Hydrallmania</i>	Genus/species	4, 5, 6	Hyd	
I	Infralittoral	Habitat factor	2, 3	I	
K	Kelps	Taxon group	4, 5, 6	K	
L	Littoral	Habitat factor	2, 3	L	
L	Low [salinity]	Habitat factor	4, 5, 6		
L	Low energy (wave/tide sheltered)	Habitat factor	2, 3	S	
Lag	Lagoonal (low or reduced salinity)	Habitat factor	4, 5, 6	Lag	
Lan	<i>Lanice</i>	Genus/species	4, 5, 6	Lan & Lcon	
Lcor	<i>Lithothamnion corallioides</i>	Genus/species	4, 5, 6	Lcor	
Ldig	<i>Laminaria digitata</i>	Genus/species	4, 5, 6	Ldig	
Lev	<i>Levinsenia</i>	Genus/species	4, 5, 6		
Lfas	<i>Lithothamnion fasciculatum</i>	Genus/species	4, 5, 6	Lfas	
Lg	Large (solitary) [ascidians]	Community feature	4, 5, 6	SoAs	
Lgla	<i>Lithothamnion glaciale</i>	Genus/species	4, 5, 6	Lgla	
Lhof	<i>Limnodrilus hoffmeisteri</i>	Genus/species	4, 5, 6	Lim	
Lhyp	<i>Laminaria hyperborea</i>	Genus/species	4, 5, 6	Lhyp	
Lic	Lichens	Taxon group	4, 5, 6	L	
Lim	<i>Limaria</i>	Genus/species	4, 5, 6	Lim	
Lit	<i>Littorina</i>	Genus/species	4, 5, 6		
Lkor	<i>Lagis koreni</i>	Genus/species	4, 5, 6		
Loch	<i>Laminaria ochroleuca</i>	Genus/species	4, 5, 6	Loch	
Lop	<i>Lophelia</i>	Genus/species	4, 5, 6	Lop	
Lpyg	<i>Lichaena pygmaea</i>	Genus/species	4, 5, 6	Lic	
Lsac	<i>Laminaria saccharina</i>	Genus/species	4, 5, 6	Lsac	
Lum	<i>Lumbrinereis</i>	Genus/species	4, 5, 6		
M	Mid [estuarine]	Habitat factor	4, 5, 6		
M	Moderate energy (Moderately wave/tide exposed)	Habitat factor	2, 3	M	
Mac	<i>Macoma</i>	Genus/species	4, 5, 6	Mac	

Mag	<i>Magelona</i>	Genus/species	4, 5, 6	Mag	
Mal	Maldanid polychaetes	Taxon group	4, 5, 6		
Mas	<i>Mastocarpus</i>	Genus/species	4, 5, 6	Mas	
Max	<i>Maxmuelleria</i>	Genus/species	4, 5, 6		
Mdis	<i>Musculus discors</i>	Genus/species	4, 5, 6	Mus	
Med	<i>Mediomastus</i>	Genus/species	4, 5, 6		
Meg	Megafauna (burrowing)	Community feature	4, 5, 6	Meg	
Mel	<i>Mellina</i>	Genus/species	4, 5, 6		
Mo	Mobile	Habitat factor	4, 5, 6	Mob	
Mod	<i>Modiolus</i>	Genus/species	4, 5, 6	Mod	
Moe	<i>Moerella</i>	Genus/species	4, 5, 6		
Mol	<i>Molgula</i>	Genus/species	4, 5, 6	Mol	
Mp	Macrophytes (angiosperms or seaweeds)	Taxon group	2, 3		
Mrl	Maerl	Taxon group	4, 5, 6	Mrl	
Msen	<i>Metridium senile</i>	Genus/species	4, 5, 6	Met	
Msim	<i>Microphthalmus similis</i>	Genus/species	4, 5, 6		
Mu	Mud/muddy [sand]	Habitat factor	2, 3	MU	
Mus	Mussels	Taxon group	4, 5, 6	M	
MuSa	Muddy sand	Habitat factor	4, 5, 6	MS	
Mx	Mixed sediments (mixtures of gravel, sand & mud, often with shell, pebble & cobble)	Habitat factor	2, 3, 4, 5, 6	MX & Mx	
Myr	<i>Myrtea</i>	Genus/species	4, 5, 6		
Mys	<i>Mysella</i>	Genus/species	4, 5, 6		
Myt	<i>Mytilus</i>	Genus/species	4, 5, 6	Myt	
Ncir	<i>Nephtys cirrosa</i>	Genus/species	4, 5, 6	Ncir	
Nem	<i>Nemertesia</i>	Genus/species	4, 5, 6	Nem	
Neo	<i>Neocrania</i>	Genus/species	4, 5, 6	Neo	
Nhom	<i>Nephtys hombergii</i>	Genus/species	4, 5, 6	Nhom	
Nint	<i>Neomysis integer</i>	Genus/species	4, 5, 6	Neo	
Nmix	<i>Neopentadactyla mixta</i>	Genus/species	4, 5, 6	Neo	
Nten	<i>Nuculoma tenuis</i>	Genus/species	4, 5, 6		
Nuc	<i>Nucula</i>	Genus/species	4, 5, 6	Nuc	
O	Offshore cirralittoral	Habitat factor	2, 3	CO	
Obor	<i>Ophelia borealis</i>	Genus/species	4, 5, 6		
Ocn	<i>Ocnus</i>	Genus/species	4, 5, 6	Ocn	
Odub	<i>Ophryotrocha dubia</i>	Genus/species	4, 5, 6		
Ofus	<i>Owenia fusiformis</i>	Genus/species	4, 5, 6		
Ol	Oligochaetes	Taxon group	4, 5, 6	Ol	
Oph	<i>Ophiura</i>	Genus/species	4, 5, 6	Oph	
Osm	<i>Osmundea</i>	Genus/species	4, 5, 6	Osm	
Ost	<i>Ostrea</i>	Genus/species	4, 5, 6	Ost	

Ov	Overhangs	Habitat factor	4, 5, 6	Ov	
Pal	<i>Palmaria</i>	Genus/species	4, 5, 6	Pal	
Par	<i>Paracentrotus</i>	Genus/species	4, 5, 6	Par	
Paur	<i>Polyclinum aurantium</i>	Genus/species	4, 5, 6	Paur	
Pb	Pebbles	Habitat factor	4, 5, 6		
Pcal	<i>Phymatolithon calcareum</i>	Genus/species	4, 5, 6	Phy	
Pcom	<i>Porella compressa</i>	Genus/species	4, 5, 6		
Pcri	<i>Phyllophora crispa</i>	Genus/species	4, 5, 6	Pcri	
Pec	<i>Pectenogammarus</i>	Genus/species	4, 5, 6	Pec	
Pel	<i>Pelvetia</i>	Genus/species	4, 5, 6	Pel	
Pen	<i>Pentapora</i>	Genus/species	4, 5, 6		
Pful	<i>Paraonis fulgens</i>	Genus/species	4, 5, 6		
Pha	<i>Phakellia</i>	Genus/species	4, 5, 6	Pha	
Phi	<i>Philine</i>	Genus/species	4, 5, 6	Phi	
Phy	<i>Phyllophora</i>	Genus/species	4, 5, 6	Phy	
Pid	Piddocks (bivalves)	Taxon group	4, 5, 6	Pid	
Pil	<i>Pilinia</i>	Genus/species	4, 5, 6		
Pjef	<i>Paramphinome jeffreysii</i>	Genus/species	4, 5, 6		
Pk	Park (kelp)	Community feature	4, 5, 6	Pk	
Pkef	<i>Protodorvillea kefersteini</i>	Genus/species	4, 5, 6		
Plon	<i>Photis longicaudata</i>	Genus/species	4, 5, 6		
Pmax	<i>Pecten maximus</i>	Genus/species	4, 5, 6		
Pnk	Plankton	Taxon group	2, 3		
Po	Polychaetes	Taxon group	4, 5, 6	P	
Pol	<i>Polydora</i>	Genus/species	4, 5, 6	Pol	
Pom	<i>Pomatoceros</i>	Genus/species	4, 5, 6	Pom	
Pon	<i>Pontocrates</i>	Genus/species	4, 5, 6	Pon	
Por	<i>Porphyra</i>	Genus/species	4, 5, 6	Por	
Pova	<i>Parvecardium ovale</i>	Genus/species	4, 5, 6	Par	
Ppel	<i>Phaxus pellucidus</i>	Genus/species	4, 5, 6		
Pra	<i>Prasiola</i>	Genus/species	4, 5, 6	Pra	
Pro	<i>Protanthea</i>	Genus/species	4, 5, 6	Pro	
Prot	<i>Polyides rotundus</i>	Genus/species	4, 5, 6	Pol	
Psa	<i>Psammechinus</i>	Genus/species	4, 5, 6	Psa	
Pse	<i>Pseudamussium</i>	Genus/species	4, 5, 6		
Puly	<i>Patella ulyssiponensis</i>	Genus/species	4, 5, 6		
R	Red seaweeds (Rhodophyceae)	Taxon group	4, 5, 6	R	
R	Reduced [salinity]	Habitat factor	4, 5, 6	RS	
R	Reef (biogenic)	Habitat factor	2, 3		
R	Rock (bedrock, boulders, stable cobbles & pebbles)	Habitat factor	2, 3	R	
Rho	<i>Rhodothamniella</i>	Genus/species	4, 5, 6	Rho	
Rkp	Rockpools	Habitat factor	4, 5, 6	Rkp	

Rup	<i>Ruppia</i>	Genus/species	4, 5, 6	Rup	
S	Salinity (Full, Variable, Reduced, Low)	Habitat factor	4, 5, 6	S	
S	Sediment	Habitat factor	2, 3	S	
S	Sublittoral	Habitat factor	2, 3, 4, 5, 6	S	
S	Surge [gully]	Habitat factor	4, 5, 6	SG	
Sa	Sands/sandy [mud]	Habitat factor	2, 3, 4, 5, 6	Snd & S	
Sab	<i>Sabellaria</i>	Genus/species	4, 5, 6	Sab	
Sac	<i>Saccorhiza</i>	Genus/species	4, 5, 6	Sac	
Sag	<i>Sagartia</i>	Genus/species	4, 5, 6		
Salv	<i>Sabellaria alveolata</i>	Genus/species	4, 5, 6	Salv	
SaMu	Sandy mud	Habitat factor	4, 5, 6	SMu	
Sar	<i>Sargassum</i>	Genus/species	4, 5, 6	Sar	
Sco	<i>Scolecopsis</i>	Genus/species	4, 5, 6		
Scr	Scoured	Habitat factor	4, 5, 6	Scr	
Scr	<i>Scrobicularia plana</i>	Genus/species	4, 5, 6		
Scup	<i>Sertularella cupressina</i>	Genus/species	4, 5, 6	Scup	
Sec	<i>Securiflustra</i>	Genus/species	4, 5, 6	Sec	
Sed	Sediment	Habitat factor	4, 5, 6	Sed	
Sem	<i>Semibalanus</i>	Genus/species	4, 5, 6	Sem	
Ser	<i>Serpula</i>	Genus/species	4, 5, 6	Ser	
Sf	Soft [rock]	Habitat factor	4, 5, 6	SfR	
Sgr	Seagrass	Taxon group	4, 5, 6	Sgr	
Sh	Shingle	Habitat factor	4, 5, 6	Sh	
Sm	Saltmarsh	Taxon group	4, 5, 6	Sm	
Sm	Small (solitary) [ascidians]	Community feature	4, 5, 6		
Sp	Sponges	Taxon group	4, 5, 6	S	
Spav	<i>Sabella pavonina</i>	Genus/species	4, 5, 6		
Spn	Seapens	Taxon group	4, 5, 6	Sp	
Sspi	<i>Sabellaria spinulosa</i>	Genus/species	4, 5, 6	Sspi	
St	Strandline	Habitat factor	4, 5, 6		
Str	<i>Streblospio</i>	Genus/species	4, 5, 6		
Sty	<i>Styela</i>	Genus/species	4, 5, 6	Sty	
Sub	<i>Suberites</i>	Genus/species	4, 5, 6	Sub	
Sund	<i>Sagartiageton undatum</i>	Genus/species	4, 5, 6		
Sw	Seaweeds	Taxon group	4, 5, 6	Sw	
Swi	<i>Swiftia</i>	Genus/species	4, 5, 6	Swi	
T	Tide-swept	Habitat factor	4, 5, 6	T	
Tal	Talitrid amphipods	Taxon group	4, 5, 6	Tal	
Tb	Tube/tube-building	Community feature	4, 5, 6	Tube	
Tben	<i>Tubificoides benedii</i>	Genus/species	4, 5, 6	Tub	

Tf	Turf	Community feature	4, 5, 6	Tf	
Thy	<i>Thyasira</i>	Genus/species	4, 5, 6	Thy	
Tra	<i>Trailliella</i>	Genus/species	4, 5, 6	Tra	
Ttub	<i>Tubifex tubifex</i>	Genus/species	4, 5, 6	Ttub	
Tub	<i>Tubularia indivisa</i>	Genus/species	4, 5, 6	Tub	
Tubi	<i>Tubificoides</i>	Genus/species	4, 5, 6	Tub	
U	Upper [estuarine]	Habitat factor	4, 5, 6		
Ulo	<i>Ulothrix</i>	Genus/species	4, 5, 6	Ulo	
Uro	<i>Urospora</i>	Genus/species	4, 5, 6	Uro	
Urt	<i>Urticina</i>	Genus/species	4, 5, 6	Urt	
V	Variable [salinity]	Habitat factor	4, 5, 6	VS	
Ven	Venerid bivalves	Taxon group	4, 5, 6	Ven	
Ver	<i>Verrucaria</i>	Genus/species	4, 5, 6	Ver	
Vir	<i>Virgularia</i>	Genus/species	4, 5, 6	Vir	
Vmuc	<i>Verrucaria mucosa</i>	Genus/species	4, 5, 6		
Vsen	<i>Venerupis senegalensis</i>	Genus/species	4, 5, 6	Vsen	
Vt	Vertical	Habitat factor	4, 5, 6	V	
WC	Water column	Habitat factor	2, 3		
	Mixed (rocky) substrata (boulders, stones & sediment mixtures)				
X		Habitat factor	4, 5, 6	X	
XFa	Mixed fauna	Taxon group	4, 5, 6	XFa	
	Mixed foliose red seaweeds				
XFoR		Taxon group	4, 5, 6		
XK	Mixed kelps	Taxon group	4, 5, 6	XK	
YG	Yellow & grey lichens	Taxon group	4, 5, 6	YG	
Zmar	<i>Zostera marina</i>	Genus/species	4, 5, 6	Zmar	
Znol	<i>Zostera noltii</i>	Genus/species	4, 5, 6	Znol	
	<i>Abietinaria abietina</i>	Genus/species	4, 5, 6	Abi	
	Calcareous	Community feature	4, 5, 6	Ca	
	<i>Catenella caespitosa</i>	Genus/species	4, 5, 6	Cat	
	<i>Ciocalypa penicillus</i>	Genus/species	4, 5, 6	Cio	
	<i>Corbula gibba</i>	Genus/species	4, 5, 6	Cor	
	Crustose bryozoans	Taxon group	4, 5, 6	ByC	
	<i>Echinus esculentus</i>	Genus/species	4, 5, 6	Ech	
	Exposed	Habitat factor	2, 3	E	Replaced by H (high energy)
	<i>Gammarus</i>	Genus/species	4, 5, 6	Gam	
	Gravels and clean sands	Habitat factor	2, 3	GS	Replaced by CS (coarse sediments)
	<i>Halichondria panicea</i>	Genus/species	4, 5, 6	Hpan	
	<i>Lithothamnion dentatum</i>	Genus/species	4, 5, 6	Lden	
	<i>Littorina littorea</i>	Genus/species	4, 5, 6	Llit	

	Marine	Habitat factor	4, 5, 6	Mar	Replaced by FS (full salinity)
	Massive [sponges]	Community feature	4, 5, 6	MaS	
	Moderately exposed	Habitat factor	2, 3	M	Replaced by M (moderate energy)
	<i>Mya arenaria</i>	Genus/species	4, 5, 6	Mare	
	<i>Mya truncata</i>	Genus/species	4, 5, 6	Mtru	
	<i>Ophiopholis aculeata</i>	Genus/species	4, 5, 6	Oacu	
	Oysters	Taxon group	4, 5, 6	Oy	
	<i>Parasmittina trispinosa</i>	Genus/species	4, 5, 6	Par	
	<i>Patella</i>	Genus/species	4, 5, 6	Pat	
	<i>Polycarpa</i>	Genus/species	4, 5, 6	Pol	
	<i>Polymastia boletiformis</i>	Genus/species	4, 5, 6	Pbol	
	<i>Pygospio elegans</i>	Genus/species	4, 5, 6	Pyg	
	<i>Sertularia argentea</i>	Genus/species	4, 5, 6	Ser	
	Sheltered	Habitat factor	2, 3	S	Replaced by L (low energy)
	<i>Spisula elliptica</i>	Genus/species	4, 5, 6	Sell	
	<i>Stolonica</i>	Genus/species	4, 5, 6	Sto	
	Synaptid holothurians	Taxon group	4, 5, 6	Syn	
	Tubeworms	Taxon group	4, 5, 6	Tw	
	<i>Zostera</i>	Genus/species	4, 5, 6	Zos	

Field recording and data management

MNCR field recording techniques are described in Hiscock (1996), with Appendix 8 providing the guidance on how to complete MNCR field recording forms (the forms can be downloaded from the JNCC website, www.jncc.gov.uk).

Procedural Guidelines for a wide range of field sampling techniques are given in the Marine Monitoring Handbook (Davies *et al.* 2001, also available to download from the JNCC website).

The Marine Recorder database application has been specifically developed to accept marine biological data from a wide range of survey techniques, including the data held originally in the MNCR database. The application can be downloaded from the JNCC website, and includes a dictionary of the habitat classification types.

Terms used for field recording and habitat definition

The following definitions are taken from guidance notes for MNCR field recording (Appendix 8 in Hiscock *ed.* 1996). Some terms are modified for use in the classification.

Salinity - The categories are defined as follows (the points of separation approximate to critical tolerance limits for marine species):

Fully marine	30-40 ‰
Variable	18-40 ‰
Reduced	18-30 ‰
Low	<18 ‰

Wave exposure - These categories take account of the **aspect** of the coast (related to direction of prevailing or strong winds), the **fetch** (distance to nearest land), its **openness** (the degree of open water offshore) and its **profile** (the depth profile of water adjacent to the coast). Estimation of wave exposure requires inspection of charts and maps.

Extremely exposed	This category is for the few open coastlines which face into prevailing wind and receive oceanic swell without any offshore breaks (such as islands or shallows) for several thousand km and where deep water is close to the shore (50 m depth contour within about 300 m, e.g. Rockall).
Very exposed	These are open coasts which face into prevailing winds and receive oceanic swell without any offshore breaks (such as islands or shallows) for several hundred km but where deep water is not close (>300 m) to the shore. They can be adjacent to extremely exposed sites but face away from prevailing winds (here swell and wave action will refract towards these shores) or where, although facing away from prevailing winds, strong winds and swell often occur (for instance, the east coast of Fair Isle).
Exposed	At these sites, prevailing wind is onshore although there is a degree of shelter because of extensive shallow areas offshore, offshore obstructions, a restricted (<90°) window to open water. These sites will not generally be exposed to strong or regular swell. This can also include open coasts facing away from prevailing winds but where strong winds with a long fetch are frequent.
Moderately exposed	These sites generally include open coasts facing away from prevailing winds and without a long fetch but where strong winds can be frequent.

<i>Sheltered</i>	At these sites, there is a restricted fetch and/or open water window. Coasts can face prevailing winds but with a short fetch (say <20 km) or extensive shallow areas offshore or may face away from prevailing winds.
<i>Very sheltered</i>	These sites are unlikely to have a fetch greater than 20 km (the exception being through a narrow (<30°) open water window, they face away from prevailing winds or have obstructions, such as reefs, offshore.
<i>Extremely sheltered</i>	These sites are fully enclosed with fetch no greater than about 3 km.
<i>Ultra sheltered</i>	Sites with fetch of a few tens or at most 100s of metres.

In the habitat classification **exposed** (as in **exposed littoral rock**) encompasses the **extremely exposed**, **very exposed** and **exposed** categories, whilst **sheltered** (as in **sheltered littoral rock**) encompasses **sheltered** to **ultra sheltered** categories.

Tidal currents (or streams) (maximum at surface) - This is maximum tidal current strength which affects the actual area surveyed. **Note for shores and inshore areas this may differ considerably from the tidal currents present offshore.** In some narrows and sounds the top of the shore may only be covered at slack water, but the lower shore is subject to fast running water.

<i>Very strong</i>	>6 knots	(>3 m/sec.)
<i>Strong</i>	3-6 knots	(>1.5-3 m/sec.)
<i>Moderately strong</i>	1-3 knots	(0.5-1.5 m/sec.)
<i>Weak</i>	<1 knot	(<0.5 m/sec.)
<i>Very weak</i>	Negligible	

In the habitat classification tide-swept habitats typically have moderately strong or stronger tidal currents.

Zone - These definitions primarily relate to rocky habitats or those where algae grow (e.g. stable shallow sublittoral sediments). For use of the terms **infralittoral** and **circalittoral** in the classification, especially for sediments, refer also to Table 5.

<i>Supralittoral</i>	Colonised by yellow and grey lichens, above the <i>Littorina</i> populations but generally below flowering plants.
<i>Upper littoral fringe</i>	This is the splash zone above High Water of Spring Tides with a dense band of the black lichen by <i>Verrucaria maura</i> . <i>Littorina saxatilis</i> and <i>Littorina neritoides</i> often present. May include saltmarsh species on shale/pebbles in shelter.
<i>Lower littoral fringe</i>	The <i>Pelvetia</i> (in shelter) or <i>Porphyra</i> (exposed) belt. With patchy <i>Verrucaria maura</i> , <i>Verrucaria mucosa</i> and <i>Lichina pygmaea</i> present above the main barnacle population. May also include saltmarsh species on shale/pebbles in shelter.
<i>Upper eulittoral</i>	Barnacles and limpets present in quantity or with dense <i>Fucus spiralis</i> in sheltered locations.
<i>Mid eulittoral</i>	Barnacle-limpet dominated, sometimes mussels or dominated by <i>Fucus vesiculosus</i> and <i>Ascophyllum nodosum</i> in sheltered locations. <i>Mastocarpus stellatus</i> and <i>Palmaria palmata</i> patchy in lower part. Usually quite a wide belt.
<i>Lower eulittoral</i>	<i>Fucus serratus</i> , <i>Mastocarpus stellatus</i> , <i>Himantalia elongata</i> or <i>Palmaria palmata</i> variously dominant; barnacles sparse.

<i>Sublittoral fringe</i>	Dominated by <i>Alaria esculenta</i> (very exposed), <i>Laminaria digitata</i> (exposed to sheltered) or <i>Laminaria saccharina</i> (very sheltered) with encrusting coralline algae; barnacles sparse.
<i>Upper infralittoral</i>	Dense forest of kelp.
<i>Lower infralittoral</i>	Sparse kelp park, dominated by foliose algae except where grazed. May lack kelp.
<i>Upper circalittoral</i>	Dominated by animals, lacking kelp but with sparse foliose algae except where grazed.
<i>Lower circalittoral</i>	Dominated by animals with no foliose algae but encrusting coralline algae.

Substratum

<i>Bedrock</i>	Includes very soft rock-types such as chalk, peat and clay.
<i>Boulders</i>	Very large (>1024 mm), large (512-1024 mm), small (256-512 mm)
<i>Cobbles</i>	64-256 mm
<i>Pebbles</i>	16-64 mm
<i>Gravel</i>	4-16 mm
<i>Coarse sand</i>	1-4 mm
<i>Medium sand</i>	0.25-1 mm
<i>Fine sand</i>	0.063 - 0.25 mm
<i>Mud</i>	<0.063 mm (the silt/clay fraction)

Each division of sediment type above represents two divisions on the Wentworth scale (Wentworth 1922).

In the habitat classification, bedrock, stable boulders, cobbles or pebbles and habitats of mixed boulder, cobble, pebble and sediment (**mixed substrata**) as well as artificial substrata (concrete, wood, metal) are collectively referred to as **rock**. Highly mobile cobbles and pebbles (shingle), together with gravel and coarse sand are collectively referred to as **coarse sediments**. **Mixed sediment** consists of heterogeneous mixtures of gravel, sand and mud and may often have shells and stones also.

MNCR SACFOR abundance scales

The MNCR cover/density scales adopted from 1990 provide a unified system for recording the abundance of marine benthic flora and fauna in biological surveys (Connor & Hiscock 1996). The scales are given below and should be used in conjunction with the following notes:

1. Whenever an attached species covers the substratum and percentage cover can be estimated, that scale should be used in preference to the density scale.
2. Use the *massive/turf* percentage cover scale for all species, excepting those given under *crust/meadow*.
3. Where two or more layers exist, for instance foliose algae overgrowing crustose algae, total percentage cover can be over 100% and abundance grades will reflect this.
4. Percentage cover of littoral species, particularly the furoid algae, must be estimated when the tide is out.
5. Use quadrats as reference frames for counting, particularly when density is borderline between two of the scale.
6. Some extrapolation of the scales may be necessary to estimate abundance for restricted habitats such as rockpools.
7. The species (as listed over) take precedence over their actual size in deciding which scale to use.
8. When species (such as those associated with algae, hydroid and bryozoan turf or on rocks and shells) are incidentally collected (i.e. collected with other species that were specifically collected for identification) and no meaningful abundance can be assigned to them, they should be noted as present (P).

MNCR SACFOR abundance scales

S = Superabundant, A = Abundant, C = Common, F = Frequent, O = Occasional, R = Rare

GROWTH FORM			SIZE OF INDIVIDUALS / COLONIES				DENSITY
% COVER	CRUST / MEADOW	MASSIVE / TURF	<1 cm	1-3 cm	3-15 cm	>15 cm	
>80%	S		S				>1 / 0.0001 m ² (1x1 cm) >10,000 / m ²
40-79%	A	S	A	S			1-9 / 0.001 m ² (3.16x3.16 cm) 1000-9999 / m ²
20-39%	C	A	C	A	S		1-9 / 0.01 m ² (10x10 cm) 100-999 / m ²
10-19%	F	C	F	C	A	S	1-9 / 0.1 m ² 10-99 / m ²
5-9%	O	F	O	F	C	A	1-9 / m ²
1-5% or density	R	O	R	O	F	C	1-9 / 10 m ² (3.16x3.16 m)
<1% or density		R		R	O	F	1-9 / 100 m ² (10x10 m)
					R	O	1-9 / 1000 m ² (31.6x31.6 m)
						R	>1 / 10,000 m ² (100x100 m) <1 / 1000 m ²

PORIFERA	Crusts <i>Halichondria</i>	Massive spp. <i>Pachymatisma</i>		Small solitary <i>Grantia</i>	Large solitary <i>Stelligera</i>
HYDROZOA		Turf species <i>Tubularia</i> <i>Abietinaria</i>		Small clumps <i>Sarsia</i> <i>Aglaophenia</i>	Solitary <i>Corymorpha</i> <i>Nemertesia</i>
ANTHOZOA	<i>Corynactis</i>	<i>Alcyonium</i>		Small solitary <i>Epizoanthus</i> <i>Caryophyllia</i>	Med. Solitary <i>Virgularia</i> <i>Cerianthus</i> <i>Urticina</i> Large solitary <i>Eunicella</i> <i>Funiculina</i> <i>Pachycerianthus</i>
ANNELIDA	<i>Sabellaria spinulosa</i>	<i>Sabellaria alveolata</i>	<i>Spirorbis</i>	Scale worms <i>Nephtys</i> <i>Pomatoceros</i>	<i>Chaetopterus</i> <i>Arenicola</i> <i>Sabella</i>
CRUSTACEA	Barnacles Tubicolous amphipods		<i>Semibalanus</i> Amphipods	<i>B. balanus</i> <i>Anapagurus</i> <i>Pisidia</i>	<i>Pagurus</i> <i>Galathea</i> Small crabs <i>Hyas araneus</i> <i>Homarus</i> <i>Nephrops</i>
MOLLUSCA			Small gastropod <i>L. neritoides</i>	Med. gastropod <i>L. littorea</i> <i>Patella</i>	Large gastropod <i>Buccinum</i> Lge bivalves <i>Mya, Pecten</i> <i>Arctica</i>
	<i>Mytilus</i> <i>Modiolus</i>		Small bivalves <i>Nucula</i>	Med. bivalves <i>Mytilus</i> <i>Pododesmus</i>	
BRACHIOPODA				<i>Neocrania</i>	
BRYOZOA	Crusts	<i>Pentapora</i> <i>Bugula Flustra</i>			<i>Alcyonidium</i> <i>Porella</i>
ECHINO- DERMATA				<i>Antedon</i> Small starfish <i>Echinocyamus</i> <i>Ocnus</i>	Large starfish <i>Echinus</i> <i>Echinocardium</i> <i>Aslia, Thyone</i> <i>Holothuria</i>
ASCIDIACEA	Colonial <i>Dendrodoa</i>			Small solitary <i>Dendrodoa</i>	Large solitary <i>Ascidia, Ciona</i> <i>Diazona</i>
PISCES				Gobies Blennies	Dog fish Wrasse
PLANTS	Crusts, Maerl <i>Audouinella</i> Fucoids, Kelp <i>Desmarestia</i>	Foliose Filamentous		<i>Zostera</i>	Kelp <i>Halidrys</i> <i>Chorda</i> <i>Himanthalia</i>

Examples of groups or species for each category

References

- Clarke, K.R., & Warwick, R.M. 2001. *Changes in marine communities: an approach to statistical analysis and interpretation*. 2nd ed. PRIMER-E. Plymouth.
- Connor, D.W., Hill, T.O., Little, M.C., & Northen, K.O. 1995a. Marine Nature Conservation Review: intertidal biotope manual. Version 6.95. *JNCC Report*, No. 249.
- Connor, D.W., Hiscock, K., Foster-Smith, R.L., & Covey, R. 1995b. A classification system for benthic marine biotopes. In: *Biology and ecology of shallow coastal waters. Proceedings of the 28th European Marine Biology Symposium*, ed. by A. Eleftheriou, A.D. Ansell & C.J. Smith, 155-165. Fredensborg, Olsen & Olsen.
- Connor, D.W., Brazier, D.P., Hill, T.O., Holt, R.H.F., Northen, K.O., & Sanderson, W.G. 1996. *Marine Nature Conservation Review: marine biotopes. A working classification for the British Isles*. Version 96.7. Peterborough, Joint Nature Conservation Committee.
- Connor, D.W., & Hiscock, K. 1996. Data collection methods (with Appendices 5 - 10). In: *Marine Nature Conservation Review: rationale and methods*, ed. by K. Hiscock, 51-65, 126-158. Peterborough, Joint Nature Conservation Committee. (Coasts and seas of the United Kingdom. MNCR series.)
- Connor, D.W. ed. 1997. *Classification of benthic marine biotopes of the north-east Atlantic. Proceedings of the second BioMar-Life workshop, Dublin, 10 September 1995*. Unpublished, Joint Nature Conservation Committee.
- Connor, D.W., Brazier, D.P., Hill, T.O., & Northen, K.O. 1997a. Marine Nature Conservation Review: marine biotope classification for Britain and Ireland. Volume 1. Littoral biotopes. Version 97.06. *JNCC Report*, No. 229.
- Connor, D.W., Dalkin, M.J., Hill, T.O., Holt, R.H.F., & Sanderson, W.G. 1997b. Marine Nature Conservation Review: marine biotope classification for Britain and Ireland. Volume 2. Sublittoral biotopes. Version 97.06. *JNCC Report*, No. 230.
- Connor, D.W., Breen, J., Champion, A., Gilliland, P.M., Huggett, D., Johnston, C., Laffoley, D. d'A., Lieberknecht, L., Lumb, C., Ramsay, K., and Shardlow, M. 2002. *Rationale and criteria for the identification of nationally important marine nature conservation features and areas in the UK*. Version 02.11. Peterborough, Joint Nature Conservation Committee (on behalf of the statutory nature conservation agencies and Wildlife and Countryside Link) for the Defra Working Group on the Review of Marine Nature Conservation.
- Day, J. & Roff, J. 2000. *Planning for representative marine protected areas. A framework for Canada's oceans and the Great Lakes*. Toronto, Worldwide Fund for Nature (Canada).
- Davies, J., Baxter, J., Bradley, M., Connor, D., Khan, J., Murray, E., Sanderson, W., Turnbull, C., & Vincent, M. 2001. *Marine monitoring handbook*. Peterborough, Joint Nature Conservation Committee (UK Marine SACs Project).
- Defra. 2002. *Safeguarding our seas. A strategy for the conservation and sustainable development of our marine environment*. London, Department for Environment, Food and Rural Affairs.
- Doody, J.P., Johnston, C., & Smith, B. eds. 1993. *Directory of the North Sea coastal margin*. Peterborough, Joint Nature Conservation Committee.
- Folk, R. L. 1954. The distinction between grain size and mineral composition in sedimentary nomenclature. *Journal of Geology*, **62**, 344-359.
- Guiry, M.D. & Dhooncha, E. N. 2002. AlgaeBase. World-wide Web electronic publication www.algaebase.org [6 February 2003]).

- Hiscock, K., & Connor, D.W. 1991. Benthic marine habitats and communities in Great Britain: the development of an MNCR classification. *JNCC Report*, No. 6. (Marine Nature Conservation Review Report, No. MNCR/OR/14.)
- Hiscock, K. ed. 1995. *Classification of benthic marine biotopes of the north-east Atlantic. Proceedings of a BioMar-Life workshop held in Cambridge. 16-18 November 1994*. Peterborough, Joint Nature Conservation Committee.
- Hiscock, K. ed. 1996. *Marine Nature Conservation Review: rationale and methods*. Peterborough, Joint Nature Conservation Committee. (Coasts and seas of the United Kingdom. MNCR series.)
- Howson, C.M., & Picton, B.E. eds. 1997. *The species directory of the marine fauna and flora of the British Isles and surrounding seas*. Belfast, Ulster Museum and Ross-on-Wye, Marine Conservation Society.
- <http://mrw.wallonie.be/dgrne/sibw/EUNIS/home.html>
- Joint Nature Conservation Committee. 1996. *Guidelines for selection of biological SSSIs: intertidal marine habitats and saline lagoons*. Peterborough, Joint Nature Conservation Committee.
- Laffoley, D. d'A., Connor, D.W., Tasker, M.L. & Bines, T. 2000. *Nationally important seascapes, habitats and species. A recommended approach to their identification, conservation and protection*. Prepared for the DETR Working Group on the Review of Marine Nature Conservation by English Nature and the Joint Nature Conservation Committee. Peterborough: *English Nature Research Reports*, No. 392. 17 pp.
- Mills, D.J.L. 1994. A manual for the analysis of data held on the Marine Nature Conservation Review database. *JNCC Report*, No. 173. (Marine Nature Conservation Review Report, No. MNCR/OR/18.)
- North Sea Conference. 2002. *Ministerial Declaration of the fifth international conference on the protection of the North Sea*. Bergen, Norway. 20–21 March 2002.
- Purvis, O.W., Coppins, B.J., Hawksworth, D.L., James, P.W., & Moore, D.M. eds. 1992. *The lichen flora of Great Britain and Ireland*. London, Natural History Museum for British Lichen Society.
- Rodwell, J.S. ed. 1995. *British plant communities. Volume 4. Aquatic communities, swamps and tall herb fens*. Cambridge, Cambridge University Press.
- Rodwell, J.S. 2000. *British plant communities. Volume 5: Maritime communities and vegetation of open habitats*. Cambridge, Cambridge University Press.
- Stace, C. (ed). 1991. *New flora of the British Isles*. 1st ed. Cambridge, Cambridge University Press.
- Wentworth, C.K. 1922. A scale of grade and class terms for clastic sediments. *Journal of Geology*, 30: 377-392.8