

Agenda Item 11.2 – Paper 28.8ii

Paper for HBDSEG Meeting - 9th-10th October 2013

Healthy and Biologically Diverse Seas Evidence Group

**Progress towards the development of a standardised UK
pressure-activities matrix**

Action: Attendees of the HBDSEG meeting are invited to:

- a) Agree and comment on the methodology used to review existing pressure-activities matrices.
- b) Provide comment on data sources used in the study in writing after the meeting.
- c) Provide comment on the pressure-activities links identified in writing after the meeting.

Paper by: April Eassom (JNCC) and Nicola Church (JNCC)

1 Introduction

Coastal and marine ecosystems have historically been subjected to a number of human-induced pressures associated with a variety of marine activities and developments. The mechanisms through which specific human activities impact the marine environment, however, are still poorly understood. This paper reviews previous attempts to link human activities to marine pressures in order to move towards development of a standardised UK pressure-activities matrix in which pressure-activities links are identified.

Below is a description of the planned process for progressing towards production of a standardised pressure-activities matrix.

2 Methods

A list of the data sources considered within this analysis is provided in:

Annex 2 - Excel workbook: Combined_P_A_Matrix (Tab 2: Data sources)

2.1 Step 1 – Agreeing a List of Set Marine Pressures

Pressure is defined as the mechanism through which an activity impacts the marine environment. Before a standardised UK pressure-activities matrix can be created a complete list of marine pressures must be defined. A list of marine pressures has been formally agreed by the OSPAR Intercessional Correspondence Group on Cumulative Effects (ICG-C). It is this ICG-C list of pressures that JNCC recommends for use within the standardised UK pressure-activities matrix.

Annex 2 - Excel workbook: Combined_P_A_Matrix (Tab 3: Pressure lists equivalency)

2.2 Step 2 – Agreeing a List of Human Activities

Numerous studies have attempted to describe human activities believed to be associated with marine pressures. To date however, no agreed list of human activities exists. Before a standardised UK pressure-activities matrix can be created a standard list of human activities occurring within the marine environment must be defined. In an attempt to identify the most appropriate list of human activities to take forward for use in the standardised UK pressure-activities matrix, this study compared multiple lists of human activities compiled from a variety of sources (listed in Annex 2 - Tab 2). An equivalency table displaying the relationships between each of the assessed human activities lists is provided in:

Annex 2 - Excel workbook: Combined_P_A_Matrix (Tab 4: Activity equivalency)

The list of human activities recommended to be taken forward in this study for inclusion within the standardised UK pressure-activities matrix is a modified version of the analytical framework currently being developed by Boteler *et al.* (in press) and is provided in:

Annex 2 - Excel workbook: Combined_P_A_Matrix (Tab 4: Columns A & B)

2.3 Step 3 – Assigning Pressure-Activity Links

Now that the human activities and marine pressures have been defined, an assessment of the potential links between activities and pressures can be undertaken.

Five out of the eleven studies presented in Annex 2 (Tab 2) conducted an analysis on the links between human activities and marine pressures and presented some form of pressure-activities matrix. These studies are:

1. Business as Usual (BaU) Study (ABPmer, 2012).
2. MarLIN, Marine and coastal activities to environmental factors matrix (MarLIN, 2003).
3. Natural England and JNCC simplified pressure-activities matrix (NE & JNCC Matrix published in the MB0102 sensitivity work).
4. Options for Developing Ecosystem Based Management (ODEMM) linkage framework (Koss *et al.*, 2011).
5. Scottish MPA Features, Activities, Sensitivities Tool (FEAST, 2013).

The pressure-activities links identified in each of these five matrices were collated and combined to produce a single matrix in which all previously identified pressure-activities links are highlighted. The combined matrix represents all links made by the data sources and does not consider why one study chose not to make a link while another did. This study does not consider the justifications behind these pressure-activities links. The group may wish to consider if that work is required and how it might be undertaken. This combined pressure-activities matrix can be viewed here:

Annex 2 - Excel workbook: Combined_P_A_Matrix (Tab 5: Combined matrix)

It should be noted that associations identified within the combined matrix in this document are likely to be constantly evolving; they are intended as a guide only.

3 Discussion

This study represents a comparison of the most up-to-date published work on pressure-activities links and identifies all links evidenced through current work streams.

4 References

ABPMER. 2012. *Business as Usual Projections of the Marine Environment, to inform the UK implementation of the Marine Strategy Framework Directive* [online]. Defra report No. ME5104. Available from: <http://randd.defra.gov.uk/Default.aspx?Menu=Menu&Module=More&Location=None&Completed=0&ProjectID=17775>

BOTELER, B., LAGO, M., PIET, G.J. & VAN DER WAL, J.T. 2013. EEA activity: Task 1.5.2.c Maritime activities, ETC/ICM task Milestone 2: Development of analytical framework of maritime sectors. **PAPER STILL IN DEVELOPMENT**

ECORYS. 2012. *Blue growth, scenarios and drivers for sustainable growth from oceans, seas and coasts* [online]. European Commission, DG MARE. Available from: http://ec.europa.eu/maritimeaffairs/documentation/studies/documents/blue_growth_third_interim_report_en.pdf

EUROPEAN COMMISSION (EC). 2012. *Guidance for 2012 reporting under the Marine Strategy Framework Directive, using the MSFD database tool (Version 1.0)* [online]. DG Environment, Brussels. Available from:
http://icm.eionet.europa.eu/schemas/dir200856ec/resources/MSFD%202012%20reporting%20guidance_incl_database_v1.0.pdf

FEAST. 2013. *Features, activities, sensitivities tool* [online]. SNH & JNCC project. Available from: www.marine.scotland.gov.uk/FEAST/

HM GOVERNMENT, 2011. *UK marine policy statement* [online]. The Stationary Office, London. ISBN: 978-0-10-851043-4. Available from:
<https://www.gov.uk/government/publications/uk-marine-policy-statement>

KOSS, R.S., KNIGHTS, A.M., ERIKSSON, A. & ROBINSON, L.A. 2011. *ODEMM linkage framework userguide* [online]. ODEMM guidance document series No. 1 EC FP7 Project 2442273. University of Liverpool. Available from:
http://www.liv.ac.uk/media/livacuk/odemmm/docs/ODEMM_Linkage_Framework.pdf

KOSS, R.S., KNIGHTS, A.M., ERIKSSON, A. & ROBINSON, L.A. 2011. *ODEMM Linkage Tables (Version 1)* [online]. ODEMM Guidance Document Series No. 1 EC FP7 project 2442273. University of Liverpool. Available from:
http://www.liv.ac.uk/media/livacuk/odemmm/images/ODEMM_Linkage_Tables2.xls

MARLIN. 2003. *Matrix linking maritime activities and environmental factors* [online]. The Marine Life Information Network. Available from:
<http://www.marlin.ac.uk/marinenaturaleffects.php>

UKMMAS (UK Marine Monitoring and Assessment Strategy Group) 2010. *Charting Progress 2: Productive Seas Evidence Group Feeder Report* [online], Defra report. Available from:
<http://chartingprogress.defra.gov.uk/resources>

Annex 1

Pressure list and descriptions

Created by the Intersessional Correspondence Group on Cumulative Effects

Presented by the United Kingdom and the Netherlands on behalf of the Intersessional Correspondence Group Cumulative Effects

This is an amended version of the document submitted to both EIHA and ICG-COBAM based on comments received from the Netherlands, Spain, Germany, France ICG-COBAM and the UK. Given the range of responses not all suggested revisions have been applied verbatim, however, it is believed that the spirit and intention of all the recommendations from Contracting Parties listed above have been included.

Comments have been added by Roy Lowry of BODC through the Valmer Project and Adrian Judd (ICG-C) has responded to these comments

Pressure theme	Pressures	Code	Pressure Descriptor	MSFD Annex III Table 2
Hydrological changes (inshore/local)	Temperature changes - local	H1	Events or activities increasing or decreasing local water temperature. This is most likely from thermal discharges, e.g. the release of cooling waters from power stations. This could also relate to temperature changes in the vicinity of operational sub sea power cables. This pressure only applies within the thermal plume generated by the pressure source. It excludes temperature changes from global warming which will be at a regional scale (and as such are addressed under the climate change pressures).	Significant changes in thermal regime (e.g. by outfalls from power stations)
Hydrological changes (inshore/local)	Salinity changes - local	H2	Events or activities increasing or decreasing local salinity. This relates to anthropogenic sources/causes that have the potential to be controlled, e.g. freshwater discharges from pipelines that reduce salinity, or brine discharges from salt cavern washings that may increase salinity. This could also include hydromorphological modification, e.g. capital navigation dredging if this alters the halodine, or erection of barrages or weirs that alter freshwater/seawater flow/exchange rates. The pressure may be temporally and spatially delineated derived from the causal event/activity and local environment.	Significant changes in salinity regime (e.g. by constructions impeding water movements, water abstraction)
Hydrological changes	Water flow (tidal current) changes –	H3	Changes in water movement associated with tidal streams (the rise and fall of the tide, riverine flows), prevailing winds and ocean currents. The pressure is therefore associated with activities	X

(inshore/local)	local, including sediment transport considerations <i>[possibly split water flow & sediment transport, i.e. separate into 'Hydrological' & 'Physical']</i>		that have the potential to modify hydrological energy flows, e.g. Tidal energy generation devices remove (convert) energy and such pressures could be manifested leeward of the device, capital dredging may deepen and widen a channel and therefore decrease the water flow, canalisation &/or structures may alter flow speed and direction; managed realignment (e.g. Wallasea, England). The pressure will be spatially delineated. The pressure extremes are a shift from a high to a low energy environment (or vice versa). The biota associated with these extremes will be markedly different as will the substrate, sediment supply/transport and associated seabed elevation changes. The potential exists for profound changes (e.g. coastal erosion/deposition) to occur at long distances from the construction itself if an important sediment transport pathway was disrupted. As such these pressures could have multiple and complex impacts associated with them.	
Hydrological changes (inshore/local)	Emergence regime changes – local, including tidal level change considerations <i>[possibly split emergence regime & tidal level changes]</i>	H4	Changes in water levels reducing the intertidal zone (and the associated/dependant habitats). The pressure relates to changes in both the spatial area and duration that intertidal species are immersed and exposed during tidal cycles (the percentage of immersion is dependant on the position or height on the shore relative to the tide). The spatial and temporal extent of the pressure will be dependant on the causal activities but can be delineated. This relates to anthropogenic causes that may directly influence the temporal and spatial extent of tidal immersion, e.g. upstream and downstream of a tidal barrage the emergence would be respectively reduced and increased, beach re-profiling could change gradients and therefore exposure times, capital dredging may change the natural tidal range, managed realignment, salt marsh creation. Such alteration may be of importance in estuaries because of their influence on tidal flushing and potential wave propagation. Changes in tidal flushing can change the sediment dynamics and may lead to changing patterns of deposition and erosion. Changes in tidal levels will only affect the emergence regime in areas that are inundated for only part of the time. The effects that tidal level changes may have on sediment transport are not restricted to these areas, so a very large construction could significantly affect the tidal level at a deep site without changing the emergence regime. Such a change could still have a serious impact. This excludes pressure from sea level rise which is considered under the climate change pressures.	X
Hydrological changes (inshore/local)	Wave exposure changes - local	H5	Local changes in wave length, height and frequency. Exposure on an open shore is dependant upon the distance of open seawater over which wind may blow to generate waves (the fetch) and the strength and incidence of winds. Anthropogenic sources of this pressure include artificial reefs, breakwaters, barrages, wrecks that can directly influence wave action or activities that may locally affect the incidence of winds, e.g. a dense network of wind turbines may have the potential to influence wave exposure, depending upon their location relative to the coastline.	X
Pollution and other chemical changes	Transition elements & organo-metal (e.g. TBT) contamination. Includes those	P1	The increase in transition elements levels compared with background concentrations, due to their input from land/riverine sources, by air or directly at sea. For marine sediments the main elements of concern are Arsenic, Cadmium, Chromium, Copper, Mercury, Nickel, Lead and Zinc. Organo-metallic compounds such as the butyl tins (Tri butyl tin and its derivatives) can be highly	Introduction of non-synthetic substances and compounds (e.g. heavy metals, hydro-carbons, resulting, for example, from

	priority substances listed in Annex II of Directive 2008/105/EC.		persistent and chronic exposure to low levels has adverse biological effects, e.g. imposex in molluscs.	pollution by ships and oil, gas and mineral exploration, atmospheric deposition, riverine inputs)
Pollution and other chemical changes	Hydrocarbon & PAH contamination. Includes those priority substances listed in Annex II of Directive 2008/105/EC.	P2	<p>Increases in the levels of these compounds compared with background concentrations. Naturally occurring compounds, complex mixtures of two basic molecular structures:</p> <ul style="list-style-type: none"> - straight chained aliphatic hydrocarbons (relatively low toxicity and susceptible to degradation) - multiple ringed aromatic hydrocarbons (higher toxicity and more resistant to degradation) <p>These fall into three categories based on source (includes both aliphatics and polycyclic aromatic hydrocarbons):</p> <ul style="list-style-type: none"> - petroleum hydrocarbons (from natural seeps, oil spills and surface water run-off) - pyrogenic hydrocarbons (from combustion of coal, woods and petroleum) - biogenic hydrocarbons (from plants & animals) <p>Ecological consequences include tainting, some are acutely toxic, carcinomas, growth defects.</p>	Introduction of non-synthetic substances and compounds (e.g. heavy metals, hydrocarbons, resulting, for example, from pollution by ships and oil, gas and mineral exploration, atmospheric deposition, riverine inputs)
Pollution and other chemical changes	Synthetic compound contamination (incl. pesticides, antifoulants, pharmaceuticals). Includes those priority substances listed in Annex II of Directive 2008/105/EC.	P3	<p>Increases in the levels of these compounds compared with background concentrations. Synthesised from a variety of industrial processes and commercial applications. Chlorinated compounds include polychlorinated biphenols (PCBs), dichlor-diphenyl-trichloroethane (DDT) & 2,3,7,8-tetrachlorodibenzo(p)dioxin (2,3,7,8-TCDD) are persistent and often very toxic. Pesticides vary greatly in structure, composition, environmental persistence and toxicity to non-target organisms. Includes: insecticides, herbicides, rodenticides & fungicides. Pharmaceuticals and Personal Care Products originate from veterinary and human applications comprising a variety of products including, Over the counter medications, fungicides, chemotherapy drugs and animal therapeutics, such as growth hormones. Due to their biologically active nature, high levels of consumption, known combined effects, and their detection in most aquatic environments they have become an emerging concern. Ecological consequences include physiological changes (e.g. growth defects, carcinomas).</p>	Introduction of synthetic compounds (e.g. priority substances under Directive 2000/60/EC which are relevant to the marine environment such as pesticides, anti-foulants, pharmaceuticals, resulting, for example, from losses from diffuse sources, pollution by ships, atmospheric deposition and biologically active substances)
Pollution and other chemical changes	Introduction of other substances (solid, liquid or gas)	P4	The 'systematic or intentional release of liquids, gases ...' (from MSFD Annex III Table 2) is being considered e.g. in relation to produced water from the oil industry. It should therefore be considered in parallel with P1, P2 and P3.	Introduction of other substances, whether solid, liquid or gas, in marine waters resulting from their systematic and/or international release into the

				marine environment, as permitted in accordance with other Community legislation and/or international conventions
Pollution and other chemical changes	Radionuclide contamination	P5	Introduction of radionuclide material, raising levels above background concentrations. Such materials can come from nuclear installation discharges, and from land or sea-based operations (e.g. oil platforms, medical sources). The disposal of radioactive material at sea is prohibited unless it fulfils exemption criteria developed by the International Atomic Energy Agency (IAEA), namely that both the following radiological criteria are satisfied: (i) the effective dose expected to be incurred by any member of the public or ships crew is 10 µSv or less in a year; (ii) the collective effective dose to the public or ships crew is not more than 1 man Sv per annum, then the material is deemed to contain de minimis levels of radioactivity and may be disposed at sea pursuant to it fulfilling all the other provisions under the Convention. The individual dose criteria are placed in perspective (i.e. very low), given that the average background dose to the UK population is ~2700 µSv/a. Ports and coastal sediments can be affected by the authorised discharge of both current and historical low-level radioactive wastes from coastal nuclear establishments.	Introduction of radio-nuclides
Pollution and other chemical changes	Nutrient enrichment	P6	Increased levels of the elements nitrogen, phosphorus, silicon (and iron) in the marine environment compared to background concentrations. Nutrients can enter marine waters by natural processes (e.g. decomposition of detritus, riverine, direct and atmospheric inputs) or anthropogenic sources (e.g. waste water runoff, terrestrial/agricultural runoff, sewage discharges, aquaculture, atmospheric deposition). Nutrients can also enter marine regions from 'upstream' locations, e.g. via tidal currents to induce enrichment in the receiving area. Nutrient enrichment may lead to eutrophication (see also organic enrichment). Adverse environmental effects include deoxygenation, algal blooms, changes in community structure of benthos and macrophytes.	Inputs of fertilisers and other nitrogen- and phosphorous-rich substances (e.g. from point and diffuse sources, including agriculture, aquaculture, atmospheric deposition)
Pollution and other chemical changes	Organic enrichment	P7	Resulting from the degraded remains of dead biota & microbiota (land & sea); faecal matter from marine animals; flocculated colloidal organic matter and the degraded remains of: sewage material, domestic wastes, industrial wastes etc. Organic matter can enter marine waters from sewage discharges, aquaculture or terrestrial/agricultural runoff. Black carbon comes from the products of incomplete combustion (PIC) of fossil fuels and vegetation. Organic enrichment may lead to eutrophication (see also nutrient enrichment). Adverse environmental effects include deoxygenation, algal blooms, changes in community structure of benthos and macrophytes.	Inputs of organic matter (e.g. sewers, mariculture, riverine inputs)
Pollution and other chemical changes	Deoxygenation	P8	Any deoxygenation that is not directly associated with nutrient or organic enrichment. The lowering, temporarily or more permanently, of oxygen levels in the water or substrate due to anthropogenic causes (some areas may naturally be deoxygenated due to stagnation of water masses, e.g. inner basins of fjords).. This is typically associated with nutrient and organic enrichment, but it can also derive from the release of ballast water or other stagnant waters (where organic or nutrient enrichment may be absent). Ballast waters may be deliberately	X

			deoxygenated via treatment with inert gases to kill non-indigenous species.	
Physical loss (Permanent Change)	Physical loss (to land or freshwater habitat)	L1	The permanent loss of marine habitats. Associated activities are land claim, new coastal defences that encroach on and move the Mean High Water Springs mark seawards, the footprint of a wind turbine on the seabed, dredging if it alters the position of the halodine. This excludes changes from one marine habitat type to another marine habitat type.	Sealing (e.g. by permanent constructions)
Physical loss (Permanent Change)	Physical change (to another seabed type)	L2	The permanent change of one marine habitat type to another marine habitat type, through the change in substratum, including to artificial (e.g. concrete). This therefore involves the permanent loss of one marine habitat type but has an equal creation of a different marine habitat type. Associated activities include the installation of infrastructure (e.g. surface of platforms or wind farm foundations, marinas, coastal defences, pipelines and cables), the placement of scour protection where soft sediment habitats are replaced by hard/coarse substrate habitats, removal of coarse substrate (marine mineral extraction) in those instances where surficial finer sediments are lost, capital dredging where the residual sedimentary habitat differs structurally from the pre-dredge state, creation of artificial reefs, mariculture i.e. mussel beds. Protection of pipes and cables using rock dumping and matting techniques. Placement of cuttings piles from oil & gas activities could fit this pressure type, however, there may be an additional pressures, e.g. "pollution and other chemical changes" theme. This pressure excludes navigation dredging where the depth of sediment is changes locally but the sediment typology is not changed.	Smothering (e.g. by man made structures, disposal of dredge spoil)
Physical damage (Reversible Change)	Habitat structure changes - removal of substratum (extraction)	D1	Unlike the "physical change" pressure type where there is a permanent change in sea bed type (e.g. sand to gravel, sediment to a hard artificial substrate) the "habitat structure change" pressure type relates to temporary and/or reversible change, e.g. from marine mineral extraction where a proportion of seabed sands or gravels are removed but a residual layer of seabed is similar to the pre-dredge structure and as such biological communities could re-colonise; navigation dredging to maintain channels where the silts or sands removed are replaced by non-anthropogenic mechanisms so the sediment typology is not changed.	Selective extraction (e.g. by exploration and exploitation of living and non-living resources on seabed and subsoil)
Physical damage (Reversible Change)	Penetration and/or disturbance of the substrate below the surface of the seabed, including abrasion	D2	The disturbance of sediments where there is limited or no loss of substrate from the system. This pressure is associated with activities such as anchoring, taking of sediment/geological cores, cone penetration tests, cable burial (ploughing or jetting), propeller wash from vessels, certain fishing activities, e.g. scallop dredging, beam trawling. Agitation dredging, where sediments are deliberately disturbed by and by gravity & hydraulic dredging where sediments are deliberately disturbed and moved by currents could also be associated with this pressure type. Compression of sediments, e.g. from the legs of a jack-up barge could also fit into this pressure type. Abrasion relates to the damage of the sea bed surface layers (typically up to 50cm depth) Activities associated with abrasion can cover relatively large spatial areas and include: fishing with towed demersal trawls (fish & shellfish); bio-prospecting such as harvesting of biogenic features such as maerl beds where, after extraction, conditions for recolonisation remain suitable or relatively localised activities including: seaweed harvesting, recreation, potting, aquaculture. Change from gravel to silt substrate would adversely affect herring spawning grounds.	Abrasion (e.g. impact on the seabed of commercial fishing, boating, anchoring)

<p>Physical damage (Reversible Change)</p>	<p>Changes in suspended solids (water clarity)</p>	<p>D3</p>	<p>Changes in water clarity from sediment & organic particulate matter concentrations. It is related to activities disturbing sediment and/or organic particulate matter and mobilising it into the water column. Could be 'natural' land run-off and riverine discharges or from anthropogenic activities such as all forms of dredging, disposal at sea, cable and pipeline burial, secondary effects of construction works, e.g. breakwaters. Particle size, hydrological energy (current speed & direction) and tidal excursion are all influencing factors on the spatial extent and temporal duration. This pressure also relates to changes in turbidity from suspended solids of organic origin (as such it exudes sediments - see the "changes in suspended sediment" pressure type). Salinity, turbulence, pH and temperature may result in flocculation of suspended organic matter. Anthropogenic sources mostly short lived and over relatively small spatial extents.</p>	<p>X</p>
<p>Physical damage (Reversible Change)</p>	<p>Siltation rate changes, including smothering (depth of vertical sediment overburden)</p>	<p>D4</p>	<p>When the natural rates of siltation are altered (increased or decreased). Siltation (or sedimentation) is the settling out of silt/sediments suspended in the water column. Activities associated with this pressure type include mariculture, land claim, navigation dredging, disposal at sea, marine mineral extraction, cable and pipeline laying and various construction activities. It can result in short lived sediment concentration gradients and the accumulation of sediments on the sea floor. This accumulation of sediments is synonymous with "light" smothering, which relates to the depth of vertical overburden.</p> <p>"Light" smothering relates to the deposition of layers of sediment on the seabed. It is associated with activities such as sea disposal of dredged materials where sediments are deliberately deposited on the sea bed. For "light" smothering most benthic biota may be able to adapt, i.e. vertically migrate through the deposited sediment.</p> <p>"Heavy" smothering also relates to the deposition of layers of sediment on the seabed but is associated with activities such as sea disposal of dredged materials where sediments are deliberately deposited on the sea bed. This accumulation of sediments relates to the depth of vertical overburden where the sediment type of the existing and deposited sediment has similar physical characteristics because, although most species of marine biota are unable to adapt, e.g. sessile organisms unable to make their way to the surface, a similar biota could, with time, re-establish. If the sediments were physically different this would fall under L2.</p> <p>Eleftheriou and Munday, 2005 describe that the majority of animals will inhabit the top 5-10 cm in open waters and the top 15 cm in intertidal areas. The depth of sediment overburden that benthic biota can tolerate is both trophic group and particle size/sediment type dependant (Bolam, 2010). Recovery from burial can occur from:</p> <ul style="list-style-type: none"> - planktonic recruitment of larvae - lateral migration of juveniles/adults - vertical migration <p>(see Chandrasekara and Frid, 1998; Bolam et al, 2003, Bolam & Whomersley, 2005). Spatial scale, timing, rate and depth of placement all contribute the relative importance of these three recovery mechanisms (Bolam et al, 2006).</p> <p>As such the terms "light" and "heavy" smothering are relative and therefore difficult to define</p>	<p>Changes in siltation (e.g. by outfalls, increased run-off, dredging/disposal or dredge spoil)</p>

			<p>in general terms. Bolam, 2010 cites various examples:</p> <ul style="list-style-type: none"> - H. ulvae maximum overburden 5 cm (Chandrasekara & Frid, 1998) - H. ulvae maximum overburden 20 cm mud or 9 cm sand (Bijerk, 1988) - S. shrebsolii maximum overburden 6 cm (Saila et al, 1972, cited by Hall 1994) - N. succinea maximum overburden 90 cm (Maurer et al 1982) - gastropod molluscs maximum overburden 15 cm (Roberts et al, 1998). <p>Bolam, 2010 also reported when organic content was low:</p> <ul style="list-style-type: none"> - H. ulvae maximum overburden 16 cm - T. benedii maximum overburden 6 cm - S. shrebsolii maximum overburden <6 cm - Tharyx sp.A. maximum overburden <6 cm 	
Other physical pressures	Litter	O1	<p>Marine litter is any manufactured or processed solid material from anthropogenic activities discarded, disposed or abandoned (excluding legitimate disposal) once it enters the marine and coastal environment including: plastics, metals, timber, rope, fishing gear etc and their degraded components, e.g. microplastic particles. Ecological effects can be physical (smothering), biological (ingestion, including uptake of microplastics; entangling; physical damage; accumulation of chemicals) and/or chemical (leaching, contamination).</p>	Marine litter
Other physical pressures	Electromagnetic changes	O2	<p>Localised electric and magnetic fields associated with operational power cables and telecommunication cables (if equipped with power relays). Such cables may generate electric and magnetic fields that could alter behaviour and migration patterns of sensitive species (e.g. sharks and rays).</p>	X
Other physical pressures	Underwater noise changes	O3	<p>Increases over and above background noise levels (consisting of environmental noise (ambient) and incidental man-made/anthropogenic noise (apparent)) at a particular location. Species known to be affected are marine mammals and fish. The theoretical zones of noise influence (Richardson et al 1995) are temporary or permanent hearing loss, discomfort & injury; response; masking and detection. In extreme cases noise pressures may lead to death. The physical or behavioural effects are dependant on a number of variables, including the sound pressure, loudness, sound exposure level and frequency. High amplitude low and mid-frequency impulsive sounds and low frequency continuous sound are of greatest concern for effects on marine mammals and fish. Some species may be responsive to the associated particle motion rather than the usual concept of noise. Noise propagation can be over large distances (tens of kilometres) but transmission losses can be attributable to factors such as water depth and sea bed topography. Noise levels associated with construction activities, such as pile-driving, are typically significantly greater than operational phases (i.e. shipping, operation of a wind farm).</p>	Underwater noise (e.g. from shipping, underwater acoustic equipment)
Other physical pressures	Introduction of light	O4	<p>Direct inputs of light from anthropogenic activities, i.e. lighting on structures during construction or operation to allow 24 hour working; new tourist facilities, e.g. promenade or pier lighting, lighting on oil & gas facilities etc. Ecological effects may be the diversion of bird species from migration routes if they are disorientated by or attracted to the lights. It is also</p>	X

			possible that continuous lighting may lead to increased algal growth.	
Other physical pressures	Barrier to species movement	O5	The physical obstruction of species movements and including local movements (within & between roosting, breeding, feeding areas) and regional/global migrations (e.g. birds, eels, salmon, whales). Both include up river movements (where tidal barrages & devices or dams could obstruct movements) or movements across open waters (offshore wind farm, wave or tidal device arrays, mariculture infrastructure or fixed fishing gears). Species affected are mostly birds, fish, mammals.	X
Other physical pressures	Death or injury by collision	O6	Injury or mortality from collisions of biota with both static &/or moving structures. Examples include: Collision with rigs (e.g. birds) or screens in intake pipes (e.g. fish at power stations) (static) or collisions with wind turbine blades, fish & mammal collisions with tidal devices and shipping (moving). Activities increasing number of vessels transiting areas, e.g. new port development or construction works will influence the scale and intensity of this pressure.	X
Biological pressures	Visual disturbance	B1	The disturbance of biota by anthropogenic activities, e.g. increased vessel movements, such as during construction phases for new infrastructure (bridges, cranes, port buildings etc), increased personnel movements, increased tourism, increased vehicular movements on shore etc disturbing bird roosting areas, seal haul out areas etc	X
Biological pressures	Genetic modification & translocation of indigenous species	B2	Genetic modification can be either deliberate (e.g. introduction of farmed individuals to the wild, GM food production) or a by-product of other activities (e.g. mutations associated with radionuclide contamination). Former related to escapees or deliberate releases e.g. cultivated species such as farmed salmon, oysters, scallops if GM practices employed. Scale of pressure compounded if GM species "captured" and translocated in ballast water. Mutated organisms from the latter could be transferred on ships hulls, in ballast water, with imports for aquaculture, a quaria, live bait, species traded as live seafood or 'natural' migration. Movement of native species to new regions can also introduce different genetic stock.	X
Biological pressures	Introduction or spread of non-indigenous species	B3	The direct or indirect introduction of non-indigenous species, e.g. chinese mitten crabs, slipper limpets, Pacific oyster and their subsequent spreading and out-competing of native species. Ballast water, hull fouling, stepping stone effects (e.g. offshore wind farms) may facilitate the spread of such species. This pressure could be associated with aquaculture, mussel or shellfishery activities due to imported seed stock imported or from accidental releases.	Introduction of non-indigenous species and translocations
Biological pressures	Introduction of microbial pathogens	B4	Untreated or insufficiently treated effluent discharges & run-off from terrestrial sources & vessels. It may also be a consequence of ballast water releases. In mussel or shellfisheries where seed stock are imported, 'infected' seed could be introduced, or it could be from accidental releases of effluvia. Escapees, e.g. farmed salmon could be infected and spread pathogens in the indigenous populations. Aquaculture could release contaminated faecal matter, from which pathogens could enter the food chain.	Introduction of microbial pathogens
Biological pressures	Removal of target species	B5	The commercial exploitation of fish & shellfish stocks, including smaller scale harvesting, angling and scientific sampling. The physical effects of fishing gear on sea bed communities are addressed by the "abrasion" pressure type D2, so B5 addresses the direct removal / harvesting of biota. Ecological consequences include the sustainability of stocks, impacting energy flows	Selective extraction of species, ... (e.g. by commercial and recreational fishing)

			through food webs and the size and age composition within fish stocks.	
Biological pressures	Removal of non-target species	B6	By-catch associated with all fishing activities. The physical effects of fishing gear on sea bed communities are addressed by the "abrasion" pressure type (D2) so B6 addresses the direct removal of individuals associated with fishing/ harvesting. . Ecological consequences include food web dependencies, population dynamics of fish, marine mammals, turtles and sea birds (including survival threats in extreme cases, e.g. Harbour Porpoise in Central and Eastern Baltic).	Selective extraction of species, including incidental non-target catches (e.g. by commercial and recreational fishing)