
BARNYARDS

C. R. Firth

OS Grid Reference: NH531470

Highlights

The sub-surface deposits at Barnyards include a sequence of estuarine sediments and peat. These provide a detailed record, supported by radiocarbon dating, of the changes in relative sea level that occurred in the Beauly Firth area during the Lateglacial and Holocene.

Introduction

The Barnyards site (NH 531470) is an area of carseland (flat, low-lying area of silty clay deposits of estuarine origin) located 0.5 km north of Beauly. The Beauly Firth area provides important evidence for reconstructing Lateglacial and Holocene sea-level changes in northern Scotland. Until recently the Beauly carseland deposits have received little detailed study. Early researchers, such as Horne and Hinxman (1914), Ogilvie (1923) and J.S. Smith (1968), noted only that Holocene raised beaches at "25 ft" and "15 ft" were present, and they made no detailed assessment of the carseland deposits. In contrast, recent studies by Sissons (1981c), Haggart (1982, 1986, 1987), Firth (1984, 1989a) and Firth and Haggart (1989) have indicated that evidence of relative sea-level changes which date back to the Loch Lomond Stadial is present within the area. From stratigraphical and morphological studies these authors have identified seven former marine levels, two of which lie buried beneath later deposits. Barnyards is a key locality for interpreting the stratigraphic evidence.

Description

Horne and Hinxman (1914) first identified raised marine deposits in the Beauly area. They reported that the village of Beauly stands on a "wide tract of marine alluvium, with a mean level of about 25 ft". Ogilvie (1923) also recognized a "25 ft" raised beach in the Beauly carselands and he suggested that a lower "15 ft" beach was present. These views were reiterated by J.S. Smith (1968), who suggested that the carseland deposits were Holocene in age. He also noted the presence of a degraded cliff landward of the carseland and he considered that this erosional feature was also of Holocene age.

Eyles and Anderson (1946) provided brief details of the deposits in the Barnyards area, recording 3.4 m of carseland clay overlying 0.3 m of peat. More detailed investigations of the stratigraphy of the Beauly Firth carselands have been undertaken subsequently by Sissons (1981c), Haggart (1982, 1986, 1987), Firth (1984) and Firth and Haggart (1989). Sissons (1981c) confined his investigations to the carselands south of the Beauly River and in the basin of Moniack Burn. He noted that the clay-silt carseland deposits are underlain by an extensive gravel layer. Sissons (1981c) indicated that the gravel layer rises landward from below -1 m OD to a maximum altitude of 2 m OD. He noted that it rests on Lateglacial marine sediments and that it terminates landward near the base of the degraded cliff that backs the carselands. Between the cliff and the buried gravel layer he identified a steeply sloping surface of erosion, which rises to 6–6.5 m OD at the base of the cliff. Sissons (1981c) concluded that these buried features were marine in origin.

Similar stratigraphical investigations were conducted by Firth (1984) on the carselands north of the Beauly River (Figure 7.20). Firth (1984) also identified a buried gravel layer rising to near 2 m OD (for example, south-east of Windhill, Figure 7.20) and a steeply sloping erosion surface which terminated at the base of a degraded cliff, at an altitude of about 8 m OD. Firth (1984) also suggested that these features were marine.




Figure 7.20: Geomorphology of the Beaully carselands (from Firth, 1984).

Haggart (1982, 1986, 1987) undertook stratigraphical investigations of the carselands between Wellhouse and Barnyards (Figure 7.20, A-B). In this area he identified a sequence of marine and terrestrial deposits (Figure 7.21). At the base of the sequence till is overlain by light-grey, silty clay. The latter is truncated by a layer of silt, sand and gravel, which is succeeded by grey, clayey silt rising to an altitude of 7.6 m OD. This layer is overlain by peat, the base of which has been dated to 9610 + 130 BP (Birm-1123). Haggart (1986) has interpreted this clayey silt/peat contact as a regressive overlap caused by a fall in relative sea level. An additional radiocarbon date from a buried peat layer farther seaward indicates that relative sea level was still falling after 9200 BP. Lying above the peat in the sequence is a grey, marine silty clay (carse deposits), which rises to a maximum altitude of 9.5 m OD. This in turn is overlain by another peat whose base has been dated to 5510+80 BP (Birm-1122) (Haggart, 1982, 1986). Haggart (1982) also identified a light-grey, micaceous silty sand within the carse deposits. He noted that this sand layer rises to a maximum altitude of 9.3 m OD and is marine in origin. A similar deposit at Moniack, 4 km to the south-east, was dated to 7270 + 90 BP (Birm-1126) (Haggart, 1982). A similar sequence of marine and terrestrial deposits was identified by Firth (1984) in the carseland north of Barnyards.

silty fine sand layer. This deposit was thought to represent either a period of increased marine transgression or a storm surge event (Haggart, 1982, 1986, 1987, 1988b). It has been correlated with similar deposits identified throughout eastern Scotland (see Silver Moss and Maryton) (Smith *et al.*, 1985a; Dawson *et al.*, 1990), and is now ascribed to a tsunami associated with the second Storegga Slide on the Norwegian continental slope (Dawson *et al.*, 1988; Long *et al.*, 1989a).

The transgressive event which deposited the coarse silty clays culminated at about 6500 BP with the formation of the highest Holocene shoreline fragment in the area, at 9.5 m OD (Haggart, 1982; Firth, 1984; Firth and Haggart, 1989). Firth (1984) correlated this shoreline fragment with the Main Postglacial Shoreline, which has been identified throughout eastern Scotland (Sissons and Smith, 1965b; J.S. Smith, 1968; Morrison *et al.*, 1981). Relative sea level then fell to its present level via intermediate shorelines at 7.5 m, 5.9 m, 4.9 m and 3.1 m OD. These shoreline fragments formed the basis of the Holocene shorelines diagram produced by Firth (1984) and Firth and Haggart (1989), which incorporated six tilted shorelines declining in altitude towards N20°E.

The sequence of deposits at Beaulieu represents a key stratigraphic record of Late Quaternary environmental and geomorphological changes in northern Scotland. In this respect it complements the interest at Munloch Valley, where the evidence is principally morphological. It demonstrates that a slow transgression during the Loch Lomond Stadial culminated in the formation of the Main Lateglacial Shoreline, and illustrates the subsequent rapid transgression that formed the marine erosion surface. The area also reveals the early Holocene regression, during which the Main Buried Beach was formed. Similarly the area records the Holocene transgression that culminated in the formation of the Main Postglacial Shoreline, as well as a possible storm surge deposit dated to 7200 BP. The four, lower, Holocene shorelines demonstrate interruptions in the subsequent fall of relative sea level to its present level.

Similar changes in relative sea level have been identified in other areas of Scotland, namely the Forth and Tay estuaries (Morrison *et al.*, 1981) and more recently at Creich in the Dornoch Firth (Smith *et al.*, 1991b). However, the Beaulieu coarse deposits are important because they contain a wealth of stratigraphical and morphological evidence within such a limited area. Haggart (unpublished data) has suggested that further stratigraphical investigations may identify an equivalent to the Lower Buried Beach of the Forth Valley (Sissons, 1966).

The area is an integral member of a national network of Quaternary sites which illustrate the changes in relative sea level in Scotland (see Silver Moss, Western Forth Valley, Carey, Dryleys, Maryton and Philorth Valley). The area also provides details which are important in the determination of regional and national patterns of isostatic movements in the British Isles (Haggart, 1989; Shennan, 1989). For example, Firth and Haggart (1989) noted apparent shifts in the isobases of different shorelines, implying, if confirmed, shifts in the centre of isostatic uplift between the formation of different shorelines (see also Gray, 1983, 1985). Furthermore, the gradients of shorelines in the Moray Firth area are steeper than expected from existing isobase maps (Sissons, 1967a, 1983a; Jardine, 1982) suggesting that the pattern of isostatic uplift in Scotland may not be represented by a simple ellipsoid (Firth and Haggart, 1989). In addition, the complexity of isostatic uplift is attested from comparison of sea-level curves from the Western Forth Valley, Lower Strathearn, the eastern Solway Firth and the inner Moray Firth, which suggests apparent differences in relative rates of uplift between the different areas (Haggart, 1989).

Conclusions

The sediments at Barnyards contain a detailed record of relative sea-level changes in the Beaulieu Firth area during the Lateglacial and Holocene (approximately the last 13,000 years). The record shows several significant fluctuations in relative sea-level position, providing valuable comparisons with the changes that are documented at reference sites elsewhere in Scotland. Together, the records from these sites allow the wider patterns of isostatic movements (movements of the Earth's crust triggered by ice-sheet growth and loading, and melting and unloading) and sea level changes to be established.

Reference list

- Cullingford, R.A., Caseldine, C.J. and Gotts, P.E. (1980) Early Flandrian land and sea-level changes in lower Strathearn. *Nature*, **184**, 159–61.
- Dawson, A.G., Long, D. and Smith, D.E. (1988) The Storegga Slides: evidence from eastern Scotland of a possible tsunami. *Marine Geology*, **82**, 271–6.
- Dawson, A.G., Smith, D.E. and Long, D. (1990) Evidence for a tsunami from a Mesolithic site in Inverness, Scotland. *Journal of Archaeological Science*, **17**, 509–12.
- Eyles, V.A. and Anderson, J.G.C. (1946) Brick clays of north-east Scotland. *Geological Survey of Britain Wartime Pamphlet*, No. 47, 89pp.
- Firth, C.R. (1984) Raised shorelines and ice limits in the inner Moray Firth and Loch Ness areas, Scotland. Unpublished PhD thesis, Coventry (Lanchester) Polytechnic.
- Firth, C.R. (1989a) Late Devensian raised shorelines and ice limits in the inner Moray Firth area, northern Scotland. *Boreas*, **18**, 5–21.
- Firth, C.R. and Haggart, B.A. (1989) Loch Lomond Stadial and Flandrian shorelines in the inner Moray Firth area, Scotland. *Journal of Quaternary Science*, **4**, 37–50.
- Gray, J.M. (1983) The measurement of shoreline altitudes in areas affected by glacio-isostasy, with particular reference to Scotland. In *Shorelines and Isostasy* (eds D.E. Smith and A.G. Dawson). London, Academic Press, 97–128.
- Gray, J.M. (1985) *Glacio-isostatic Shoreline Development in Scotland An Overview*. Department of Geography and Earth Science, Queen Mary College, University of London. Occasional Paper, No. 24, 61pp.
- Haggart, B.A. (1982) Flandrian sea-level changes in the Moray Firth area. Unpublished PhD thesis, University of Durham.
- Haggart, B.A. (1986) Relative sea-level change in the Beaully Firth Scotland. *Boreas*, **15**, 191–207.
- Haggart, B.A. (1987) Relative sea-level changes in the Moray Firth area, Scotland. In *Sea-level Changes* (eds M.J. Tooley and I. Shennan). Oxford, Blackwell, 67–108.
- Haggart, B.A. (1988b) The stratigraphy, depositional environment and dating of a possible tidal surge deposit in the Beaully Firth area, north-east Scotland. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **66**, 215–30.
- Haggart, B.A. (1989) Variations in the pattern and rate of isostatic uplift indicated by a comparison of Holocene sea-level curves from Scotland. *Journal of Quaternary Science*, **4**, 67–76.
- Horne, J. and Hinxman, L.W. (1914) The geology of the country round Beaully and Inverness: including part of the Black Isle. (Explanation of Sheet 83). Memoirs of the Geological Survey of Scotland. Edinburgh, HMSO, 108pp.
- Jardine, W.G. (1982) Sea-level changes in Scotland during the last 18,000 years. *Proceedings of the Geologists' Association*, **93**, 25–41.
- Long, D., Smith, D.E. and Dawson, A.G. (1989a) A Holocene tsunami deposit in eastern Scotland. *Journal of Quaternary Science*, **4**, 61–6.
- Morrison, J., Smith, D.E., Cullingford, R.A. and Jones, R.L. (1981) The culmination of the main postglacial transgression in the Firth of Tay area, Scotland. *Proceedings of the Geologists' Association*, **92**, 197–209.
- Ogilvie, A.G. (1923) The physiography of the Moray Firth coast. *Transactions of the Royal Society of Edinburgh*, **53**, 377–404.
- Shennan, I. (1989) Holocene crustal movements and sea-level changes in Great Britain. *Journal of Quaternary Science*, **4**, 77–89.
- Sissons, J.B. (1966) Relative sea level changes between 10,300 and 8300 BP in part of the Carse of Stirling. *Transactions of the Institute of British Geographers*, **39**, 19–29.
- Sissons, J.B. (1967a) *The Evolution of Scotland's Scenery*. Edinburgh, Oliver and Boyd, 259pp.
- Sissons, J.B. (1969) Drift stratigraphy and buried morphological features in the Grangemouth–Falkirk–Airth area, central Scotland. *Transactions of the Institute of British Geographers*, **48**, 19–50.
- Sissons, J.B. (1976a) Lateglacial marine erosion in south east Scotland. *Scottish Geographical Magazine*, **92**, 17–29.
- Sissons, J.B. (1981c) Lateglacial marine erosion and a jökulhlaup deposit in the Beaully Firth. *Scottish Journal of Geology*, **17**, 7–19.
- Sissons, J.B. (1983a) Shorelines and isostasy in Scotland. In *Shorelines and Isostasy* (eds D.E.

-
- Smith and A.G. Dawson). London, Academic Press, 209–25.
- Sissons, J.B. and Smith, D.E. (1965b) Peat bogs in a post-glacial sea and a buried raised beach in the western part of the Carse of Stirling. *Scottish Journal of Geology*, **1**, 247–55.
- Smith, D.E. (1968) Post-glacial displaced shorelines in the surface of the carse clay on the north bank of the River Forth, in Scotland. *Zeitschrift für Geomorphologie*, NF, **12**, 388–408.
- Smith, D.E., Cullingford, R.A. and Haggart, B.A. (1985a) A major coastal flood during the Holocene in eastern Scotland. *Eizeitaler und Gegenwart*, **35**, 109–18.
- Smith, D.E., Turbayne, S.C., Firth, C.R. and Brooks, C.L. (1991b) Creich. In *Late Quaternary Coastal Evolution in the Inner Moray Firth. Field Guide* (eds C.R. Firth and B.A. Haggart). London, West London Press, 33–42.