

---

# EAS NA BROIGE DEBRIS CONE

A. Werritty and L. J. McEwen

OS Grid Reference: NN192598

## Highlights

The deposits in the debris cone at Eas na Broige provide an important geomorphological and sedimentary record of slope processes during the Holocene. They show successive phases of debris-flow activity and alluvial fan development.

## Introduction

The Eas na Broige debris cone (NN 192598) is located in Glen Etive at the base of a near-vertical south-facing rock gully (Dalness Chasm) which drains Stob na Broige (956 m OD). Debris cones are fan-shaped accumulations of poorly-sorted debris formed by successive debris flows at the base of steep gullies. Such debris cones have developed extensively at the margins of valley floors in upland Scotland over the last 13,000 years and have formed in response to changes in sediment supply from adjacent gullies and slopes. Collectively these cones represent an important class of Lateglacial and Holocene landform found throughout upland Britain (Statham, 1976b; Harvey *et al.*, 1981; Innes, 1983b, 1985; Brazier *et al.*, 1988; Brazier and Ballantyne, 1989). The Eas na Broige cone is a particularly good example, and the deposits which comprise it have provided a detailed record of slope processes during the Holocene (Brazier *et al.*, 1988).

## Description

The local bedrock comprises granite on the lower slopes with rhyolite lavas on the higher ground; the Dalness Chasm having been etched from a porphyritic dyke (Bailey and Maufe, 1916). During the Loch Lomond Stadial, glacier ice extended up to 650 m OD in upper Glen Etive (Thorp, 1981a, 1986). The debris cone is thus Holocene in age.

The Eas na Broige cone comprises two units: an upper debris cone with a concave long profile and mean gradient of 13.7°, and a lower alluvial fan with a mean gradient of 6.2°. Five discrete cone and fan surfaces can be identified within the lower alluvial surfaces which are inset into the steeper debris cone surfaces. The respective volumes of the debris cone and alluvial fan have been estimated at 170,000 m<sup>3</sup> and 100 m<sup>3</sup>; the latter being entirely derived by incision and reworking of the former (Brazier, 1987).

The stratigraphy of much of the cone has been exposed by stream incision, and at two contrasting sites clearly distinguishable debris flow and fluvial sediments have been identified (Figure 10.17). At the apex of the alluvial fan (site 1) a coarse debris flow deposit lies beneath a distinctive and strongly podsolized palaeosol, radiocarbon dated at 550 ± 50 BP (SRR-2882). The overlying sediments comprise poorly-sorted alluvial gravels with only a weak soil development. Higher up the debris cone (Site 2) a second palaeosol, radiocarbon dated at 4480 ± 300 BP (SRR-2884), separates two debris-flow units. The upper of these units is the continuation of the lower unit at site 1 and it appears to be the final debris flow unit which was deposited on this part of the cone (Brazier *et al.*, 1988). Pollen samples have been collected from site 1 in order to investigate possible vegetational changes associated with the onset of fluvial reworking. An initial cover of *Corylus*, *Alnus* and *Pinus* before 550 BP was replaced by Gramineae, *Plantago* and *Calluna* after that date. A strong presence of charcoal is also recorded in the unit above the palaeosol.




Figure 10.17: Top: schematic section along the length of the Eas na Broige debris cone. Bottom: detail of section at sampling sites (from Brazier *et al.*, 1988).

## Interpretation

Three major phases in the development of this fluvially modified debris cone can be identified. The debris cone initially developed during the first 6000 years of the Holocene, with aggradation ceasing about 4000 BP as a result of exhaustion of the sediment supply through the Dalness Chasm. A prolonged period of stability then ensued until 550 BP. This was followed by a final phase in which the incision into the debris cone produced the inset alluvial fan. The pollen evidence strongly suggests that fluvial activity was contemporaneous with changes in the vegetation cover caused by human interference. The removal of the tree cover destabilized the cone surface and triggered fluvial incision. This instability has continued until the present day.

This site has a potentially wider significance in terms of it being representative of the class of fluvially modified debris cones found throughout the Highlands (Brazier *et al.*, 1988) and other parts of upland Britain (Harvey *et al.*, 1981). First, the initial accumulation of debris-flow deposits involved the reworking of sediment deposited during deglaciation. This implies that this cone, like many others in upland Britain, is "paraglacial" in origin (cf. Ryder, 1971; Church and Ryder, 1972), that is, its formation was dependent upon an abundant sediment supply following deglaciation. Once this was exhausted, aggradation on the cone ceased. Second, fluvial incision at this site is attributed to recent human disturbance of the vegetation. There are many other fluvially modified debris cones in the Scottish Highlands and upland Britain where a similar anthropogenic trigger may have initiated the same change in the process regime on debris cones (see Statham, 1976b; Harvey *et al.*, 1981; Innes, 1983b). In this latter respect the Eas na Broige cone contrasts with those in Glen Feshie, where natural processes are considered to have been responsible for reactivation (Brazier and Ballantyne, 1989).

The Eas na Broige cone is a very good example of a Holocene debris cone that has been subject to fluvial modification. Although debris cones are ubiquitous throughout the Scottish Highlands, this site in Glen Etive is unique in that the date and extent of the fluvial reworking of the original cone have been precisely determined.

The Eas na Broige debris cone provides the most detailed record currently available of Holocene sedimentation at the margins of a major valley in Scotland. Following deglaciation at approximately 10,000 BP the debris cone developed over the next 4000 years. At this time the cone surface became stabilized until about 500 years ago, when fluvial reworking of the basal part of the cone arose in response to the removal of the forest cover by human activity. The cone thus represents a particularly good example of a fluvially modified debris cone in which the most recent phase of development has been in response to human settlement on the valley floor.

## Conclusions

Eas na Broige debris cone is a fan-shaped accumulation of poorly sorted material (mixed particles of various sizes) formed by flows, from the slopes above, of rock and soil debris mixed with water. It provides an important record of slope processes during the Holocene and is representative of a type of landform and process system that occurs widely in the Highlands. In particular it shows two phases of development, the first reflecting high sediment supply following deglaciation (ice melting and retreat) and the second, the impact of forest clearance by Man. The site is not only a good landform example but has a well-documented history of development.

## Reference list

- Bailey, E.B. and Maufe, H.B. (1916) *The Geology of Ben Nevis and Glencoe, and the Surrounding Country*. (Explanation of Sheet 53). Memoirs of the Geological Survey of Scotland. Edinburgh, HMSO, 247pp.
- Brazier, V.B., Whittington, L. and Ballantyne, C.K. 1988. Holocene debris cone formation in Glen Etive, Western Grampian Highlands, Scotland. *Earth Surface Processes and Landforms*, **13**, 525–31.
- Brazier, V.B. and Ballantyne, C.K. (1989) Late Holocene debris cone evolution in Glen Feshie, western Cairngorm Mountains Scotland. *Transactions of the Royal Society of Edinburgh. Earth Sciences*, **80**, 17–24.
- Church, M. and Ryder, J.M. (1972) Paraglacial sedimentation: a consideration of fluvial processes conditioned by glaciation. *Bulletin of the Geological Society of America*, **83**, 3059–72.
- Harvey, A.M., Oldfield, F., Baron, A.F. and Pearson, G.W. (1981) Lichens, soil development and the age of Holocene valley-floor landforms: Howgill Fells, Cumbria. *Geografiska Annaler*, **66A**, 353–66.
- Innes, J.L. (1983b) Lichenometric dating of debris flow deposits in the Scottish Highlands. *Earth surface Processes and Landforms*, **8**, 579–588.
- Innes, J.L. (1985) Magnitude-frequency relations of debris flows in north-west Europe. *Geografiska Annaler*, **67A**, 23–32.
- Ryder, J.M. (1971) The stratigraphy and morphology of paraglacial alluvial fans in south-central British Columbia. *Canadian Journal of Earth Sciences*, **8**, 279–98.
- Statham, I. (1976b) Debris flows on vegetated screes in the Black Mountain, Carmarthenshire. *Earth Surface Processes*, **1**, 173–80.
- Thorp, P.W. (1981a) A trimline method for determining the upper limit of the Loch Lomond Advance glaciers: examples from the Loch Leven and Glencoe areas. *Scottish Journal of Geology*, **17**, 49–64.
- Thorp, P.W. (1986) A mountain icefield of Loch Lomond Stadial age, western Grampians, Scotland. *Boreas*, **15**, 83–97.