

Bioindicator methods for monitoring of nitrogen impacts on statutory nature conservation sites

Mark Sutton, Carole Pitcairn, Ian Leith, Netty van Dijk,
Lucy Sheppard, Sim Tang, Simon Smart, Ruth Mitchell,
David Fowler

Centre for Ecology and Hydrology


Pat Wolsely and Peter James

Natural History Museum

JNCC, EN, SNH, CCW, EHS, SEPA

Assessing air pollution impacts: Critical loads & Bioindicators

- **Critical loads & critical levels approaches**
 - Mostly national, but can be site-based.
 - Essentially only a risk assessment.
“Likelihood of change”
- **Bioindicators & biomonitoring:**
 - Actual measurement of ecosystem parameters
 - Relates directly to site “condition”
 - Can consider temporal changes at a site level



Uncertainties and needs

- Wide range of different views regarding the usefulness of bioindicators & biomonitoring for N.
- Potentially attractive for Agencies to help monitor site condition in relation to air pollution impacts and for local assessments
- Uncertainty regarding the specific benefits and limitations of the different methods.
- Need research...

JNCC & Agencies Research Project



- Stage 1:
 - Review existing methods
 - Test novel techniques at one site
 - Identify methods with high potential for further application by Agencies
- Stage 2:
 - Further test the practical application of methods recommended for application
 - Use 4 key sites for detailed methods (NH_3 , NO_x , wet deposition and controlled dose comparison)
 - Simplest methods to be then applied at UK scale.

Definitions

- Can get easily hung up on definitions – keep it simple here
- **Bioindicators:**
 - General group of approaches where biological measurement used to indicate something (e.g. might be applied at one time for spatial comparison).
- **Biomonitoring:**
 - Repeated application of bioindicator methods over time (e.g. weeks to decades)

N bioindicators: What is being indicated?

- Several purposes for N bioindicators to estimate:
 - N deposition fluxes from the atmosphere
 - Air concentrations of N species (NO_x , NH_3 etc)
 - Environmental effects of N, including physiological and environmental changes
- Should consider biomonitoring in conjunction with physical monitoring
- Can use biomonitoring results as input for local application of the critical loads approach.



Types of N bioindicator methods

Biochemical methods

- Measure the accumulation of N or a chemical/physiological response in a plant/soil component

Species composition methods

- Record the presence of certain species previously categorized according to their N preferences and generate an overall site index

Transplant methods


- Locally occurring or standardized plants are exposed to a range of N conditions and their responses assessed.

Biochemical bio-indicator methods for nitrogen

Foliar tissue N


- Most extensively studied parameter
- Response to N deposition rather than concentrations
- Uncertainties in the past particularly related to need to standardize protocols and have robust reference estimates of N deposition.

Total foliar tissue N



Results near a farm at Earlston in Scotland


Spatial variability of foliar N



Results from Leende Heide, the Netherlands

Foliar amino acids

- Large rates of accumulation
- But amino acids accumulated different between species



“Substrate N” and foliar ammonium


Substrate Nitrogen

- Total available N for growth expected to vary more than total N.
- Approximate substrate N by measuring total soluble N in leaves.
- More general than amino acids

Foliar ammonium


- Foliar ammonium represents primary pool for N compound synthesis and recycling
- Smaller pool expected to have larger response

Foliar ammonium



- Best results so far for bryophytes
- Species differences consistent with N habitat preferences
- Massive response of factor 20 over range tested.


Response of different foliar N pools



- Larger response from the smaller pool size
- Smaller pools may also respond more quickly to change in N deposition
- Foliar ammonium potentially easier to measure


Combining Bioindicators with the Critical loads approach

- Currently feasible for foliar N and foliar ammonium
- Measured biomonitor data indicates whether a site is significantly above or below the critical load



Estimating N deposition based on foliar ammonium

- Bryophytes in woodland ground flora
- Needed for more data in clean conditions to improve confidence limits




Biochemical bioindicators of plant N responses

- Response parameters include enzyme activities, soil emissions and bioassays of damage e.g. tests of frost hardiness or photosynthetic activity.
- Methods relevant to assess ecological impacts and for monitoring of general 'condition'
- But less direct than accumulation methods and more affected by other factors, so generally less well suited to assess N deposition.

Species composition bio-indicator methods for nitrogen

Ellenberg Approach for Higher Plants and Bryophytes



- Demonstrated close relationship to N deposition
- But species composition affected by many other factors, especially soil, management & light effects.
- Suited to local scale assessments and long term biomonitoring



Lichens for assessing impact of atmospheric nitrogen

- Tested:
 - several overall biodiversity measures
 - Several N indicator methods inc.
 - Ellenberg (Wirth) values
 - French (Lallmont) scale
 - Van Herk (Acidophyte / Nitrophyte classification)
 - Compared Twigs and Trunks
 - Also assessed bark pH
- Some suggestion: Lichens are particularly responding to NH_3 . Effects of NH_4^+ and NO_y are much less certain.
- Effect of NH_3 to *increase* bark pH is critical

Total Lichen biodiversity and bark pH



Sensitivity of lichens to NH_3

Acidophytes (AIW) hate NH_3




“Troll’s Beard” (*Bryoria*):
disappearing from many sites



Cladonia

Nitrophytes (NIW) love NH_3



Xanthoria
thrives by farms



Combination of Acidophytes and Nitrophytes





Ellenberg (Wirth): detailed and potential simpler system



Critical bark pH for twig and trunk acidophytes

Twig acidophytes are completely excluded above a trunk pH of 4-4.2. This equates to an NH_3 conc of around 2 ug m^{-3} .

Note outlier for Thetford trunks: rich Parmelion community (not in AIW)



Potential for two-tier Lichen approach for Nitrogen

- **Detailed method:** further develop van Herk approach, including application to twigs. Good for full site assessment by experts.
- **Simpler method:** refine simple Ellenberg approach, including application to twigs. Good for application by trained non experts to get indications at sites and for raising public awareness.

Transplant bio-indicator methods for nitrogen

Native reciprocal transplants


- Works best for bryophytes – no soil attached and more robust than lichens
- Need two or more locations with similar climate
- Measure growth rates, and N content
- Observe reduced performance in polluted conditions or recovery in clean conditions
- Good for demonstration to stakeholders to show the benefits of emission abatement at a site level.

Grass biomonitors

- *Lolium perenne* using EUROBIONET wicking system




L. perenne Biomass




Grass biomonitors

L. perenne Biomass




L. perenne above ground %N concentration




Grass biomonitors


L. perenne: Mean mg N per pot



L. perenne: Mean mg N per pot



Overview of methods tested and recommendations



Classification of method types and recommendation for further development

- **Chemical Methods**
 - Total foliar N
 - Soluble foliar N/ NH_4^+
 - Bark pH
- **Diversity Methods**
 - Ellenberg scale (Higher plants and bryophytes)
 - Van Kerk scale for lichens (inc for twigs)
 - Ellenberg (Wirth) scale for lichens (inc for twigs)
- **Transplant Methods**
 - Standardized grass transplants
 - (Native reciprocal transplants)

Conclusions

- Methods refer to different timescales – This affects the translation from bioindicating to biomonitoring.
- The methods by definition refer to an ecosystem or biological impact – therefore they give a direct implication of critical load / level exceedance
- The accumulation methods most quantitatively related to atmospheric deposition, but different different deposition components may have different effects.
- For application to statutory nature conservation sites the bioindicators can be best used for local sources with small scale transects into reserves (NO_x and NH_3 as sources).

Conclusions and next steps

- The assessment for elevated wet deposition is much less certain (due to linked altitude effects).
- The most robust assessments would use more than one method in parallel and complement physical monitoring efforts.
- Ongoing work is focusing at 4 sites for detailed methods, plus UK wide survey of simple methods.
- Contributors are invited across the Agencies to try out the simple methods next year.